

CATALOGUE OF PLANTS

Collected in GEORGIA

within 60 miles of

OKEFINOKEE SWAMP

in the counties of

Pierce, Ware, Charlton, Glynn,


Camden, Clinch ^{and} Echols

in August, 1902,

BY


ROLAND M. HARPER,

Expert, in Botanical Investigations and
Experiments, Bureau of Plant Industry,
United States Department of Agriculture.



ITINERARY

of
Roland M Harper
August, 1902



CONTENTS

1. <i>obtaining</i>	1
2. <i>of the</i>	1
3. <i>of the</i>	1
4. <i>of the</i>	1
5. <i>of the</i>	1
6. <i>of the</i>	1
7. <i>of the</i>	1
8. <i>of the</i>	1
9. <i>of the</i>	1
10. <i>of the</i>	1
11. <i>of the</i>	1
12. <i>of the</i>	1
13. <i>of the</i>	1
14. <i>of the</i>	1
15. <i>of the</i>	1
16. <i>of the</i>	1
17. <i>of the</i>	1
18. <i>of the</i>	1
19. <i>of the</i>	1
20. <i>of the</i>	1
21. <i>of the</i>	1
22. <i>of the</i>	1
23. <i>of the</i>	1
24. <i>of the</i>	1
25. <i>of the</i>	1
26. <i>of the</i>	1
27. <i>of the</i>	1
28. <i>of the</i>	1
29. <i>of the</i>	1
30. <i>of the</i>	1

AUGUST
1902

DAY OF MONTH	NUMBER	NAME OF PLANT.	HABIT IT LOCALITY ETC.	(APPROXIMATE ALTITUDE (FEET))
1	1464	<i>Planisia tenuifolia</i> T. & G.	Open places near S. F. & W. Ry. on sand-hills on left bank of Atilla River 3 miles from Waycross in Pierce Co. Perhaps introduced?	100
1	1465	<i>Warea cuneifolia</i> (Muhl.) Nutt	Do.	100
1	1466	<i>Quercus myrtifolia</i> Willd. A small tree, 6-10 ft. tall, with very numerous stout branches, fruiting abundantly. (Photo. no. 113)	Forming part of a very dense thicket on driest parts of same sand-hills, nearer the river than the preceding.	100
1	1467	<i>Hastanea peruviana</i> A shrub 6-8 ft. tall, with small thick leaves.	Do.	100
2	1468	<i>Polygala</i> sp. Small field	Rather dry pine-barrens near S. F. & W. Ry. about a mile southwest of Waycross, Ware Co.	145

AUG.	NUMBER	
2	1469	<i>Pontederia cordata</i> L.
2	1470	<i>Persea pubescens</i> (Pursh) Sargent a shrub 5-6 ft. tall
2	1470a	<i>Sphagnum Ludovicianum</i> (R. & S.) Hamet.
7 (5 a.m.)	1471	<i>Hydrocotyle umbellata</i> L.
7 5:30 a.m.)	1472	<i>Drosera intermedia</i> Layne

Low - marshy place along R.R. N.
about 4 miles south of
Hagawass, Kan. Co.

125

Moist sphagnum pine-land
about $1\frac{1}{2}$ miles southwest of
Hennmore, Mare Co.

125

Do.

125

Sphagnum bog near Bugaboo
Landing on the canal in Cheyenne
Swamp, about 7 miles west of
Camp Cornelia.

115

Do.

115

AUG.	NO.	
7	1473	<i>Sagittaria</i>
(5:40 a.m.)		
7	1473a	<i>Sphagnum Trinitense angustilimbatum</i> Varr.?
7	1474	<i>Eriophorum Virginicum</i> L. 5 ft tall. Wool nearly white.
(7:10 a.m.)		
7	1475	<i>Pieris phillyreaefolia</i> (Hook.) Oakes. An erect shrub 2-3 ft. tall.
7	1476	<i>Sarracenia minor</i> Walt. Leaves and scapes 2-3 ft. tall.

Okefenokee Swamp.

with the preceding

115

Do.

115

Sphagnum bog between Bugaboo
Island and Bugaboo Island,
west of the camp.

115

Do.

115

Do.

115

AUG.	NO.			
7	1476a	<i>Sphagnum Ludovicianum</i> R. & B. Mant.	Over the island than the preceding, in drier places.	115
7	1477	<i>Rhynchospora</i>	Do., close to edge of island.	115
7	1478	<i>Sarracenia psittacina</i> Mx.	In sphagnum nearer the landing, about the same place as no. 1476	115
7	1479	<i>Pieris phillyreaefolia</i> (Hook.) Oe. Same as no. 1475, but climbing under bark of <i>Taxodium imbricarium</i> to a height of 20-30 ft.	On bank of canal near Chase Prairie, about 10 miles from camp Cornelia.	120
7	1480	<i>Scleria trichopoda</i> G. Wright.	Very boggy place (not sphagnum) in edge of Chase Prairie.	115

Ok. not a camp

AUG

NO

7 1481 *Utricularia juncea* Val.

Keefinokee Swamp.

Growing strictly erect in dense colonies on rather firm soil (muck) in Chase Prairie, elevated an inch or so above the water.

(Photo no. 136)

7 1481a *Sphagnum Portoricense* Lamour.

In Jackson Bay, at western edge of Chase Prairie, behind second camp, 11 miles from Camp Cornelia.

8 1481b *Leucobryum glaucum* (L.) Schimp.

Do.

8 1482 *Ilex bassine* L.

Do.

8 1482a *Lycopodium* ?Do., on base of stem of *Pieris nitida*

AUG

NO

8

1482b

Odontoschisma prostratum (Sw.) PreslNear the ponding, on cypress
(*L. mexicanum*) knee.

115

8

1483

Ludwigia capitata H.
Similar to *L. capitata*, but
with inflorescence elongated
into an interrupted spike.On west bank of canal, in
Chase Prairie.

115

8

1484

Xyris Smaliana Nash.

(Photo. 136)

In small prairie about 2 miles
from Camp Cornelia.

115

(Last plant collected in the swamp)

8

1485

*Panicum*In small cypress pond (now
dry) about $\frac{1}{2}$ mile south of
Camp Cornelia, Bl. Co.

130

8

1486

Paspalum

Do

130

AUG

NC

'8 1467 *Rhynchospora distans* (Mx.) VahlIn middle of rather dry ca. by
road in pine-barrens just north
of Camp Cornelia.

135

8 1488 *Ludwigia capitata* Mx.Rather dry pine-barrens, near
same place.

135

8 1489 *Stenophyllus stenophyllus* (Ell.) Britton

Dry sandy road near same place.

140

9 1410 *Guirana scirpoides* Mx.Rather dry pine-barrens about
3 miles north of Camp Cornelia.
Abundant but scattered.

125

9 1411 *Lachnocaulon Bepidanthum*Same habitat and locality as
the preceding.

125

AUG.	NO.	
9	1492	<i>Hesperis matronalis</i> Sw. Moist pine-barrens along extinct Brunsick Pensacola RR. at midway between Camp Columbia and Folkston, Charlton Co.
11	1493	<i>Petalostemon carneus</i> L. Moistly forest floor - 9.
11	1494	<i>Eryngium Baldwinii</i> Spreng.
11	1495	<i>Juncus biflorus</i> Ell. With pretty good rootstocks
11	1496	<i>Eryngium aromaticum</i> Baldr. Just beginning to flower.

Moist pine-barrens along extinct
Brunsick Pensacola RR. at
midway between Camp Columbia
and Folkston, Charlton Co.

Rather dry pine-barrens about
 $\frac{1}{2}$ mile southwest of Folkston,
near road to Trader's Hill.

Do.

Sand-hills between Folkston
and Spanish Creek, along road
to Trader's Hill.

Do.

AUG.

NO.

11 1497 *Cracca chrysophylla* (P.) BuntzeSand is good Spanish creek,
about a mile from Trader's Hill.
(Photo no 4-1)

90

11 1498 *Aeschynomene viscidula* L.Along road through sand hills,
between Brunswick & Pensacola
R.R. and Trader's Hill. Perhaps
not indigenous?

80

11 1499 *Helix* *mexicana*
11.45
(a.m.)Bank of St. Mary's River, at
foot of bluff at Trader's Hill.

10

11 1500 *Dichromena colorata* (L.) A. S. Hitchcock
Stoloniferous

Do.

10

1501 *Hypericum*

Shrub about 4 ft. tall,

11 1501 *Hypericum multifolium*
Shrub about 4 ft. tall,
with thinner and narrower
leaves than typical *multifolium*

Do.

15

AUG	N	
11	1502	<i>Hypericum galioides</i> Lam.
11	1503	<i>Mimosa strigillosa</i> T. & G.
11	1504	<i>Froelichia floridana</i> (Walt.) Moq.
11 2:20 p.m.	1505	<i>Dichromena</i> Similar to no. 1500, but smaller, and densely and almost exclusively covering several square rods.
12	1506	<i>Melanthium disperum</i> Small. In full bloom.

With the preceding.

10

Exposed places on river bank,
perhaps introduced.

10

In sandy cultivated field
near Traders' Hill. Very
abundant. (Photo. no. 4)

80

Moist open flat place in pine-
barrens near a tributary of
Spanish Creek, along B. & P. R. R.

Sphagnum bog along a branch
near B. & P. R. R. about 2 miles
east of Holston, Charlton Co.

11

AUG	N ^o	
12	1507	<i>Habenaria integra</i> (Nutt.) Shumg. Lip crenulate, not fringed.
12	1508	<i>Habenaria blephariglotis</i> (Willd.) Torr.
12	1508a	<i>Sphagnum tenerum</i> (Ait.) Warrm.
12 (9 a.m.)	1509	<i>Gillandsia tenuifolia</i> L. 5
12 (9:54 a.m.)	1510	<i>Geucium Nashii</i> Kearney.

in the county.

(With the preceding.)

Do.

Do., toward edge of bog.

on living trunks of *Acer rubrum*
and *Nyssa biflora* in swamp
along B. & P. R.R. about 3 miles
from Folkston. (Photo. no. 140)

Low grounds in small swamp
about 4 miles from Folkston.

12

AUG

NO

12

1511

Eupatorium semiseratum DC.

(9:56

a.m.)

12

1512

Plantago sparsiflora Mx.
Like *sparsiflora*, but with
leaves thicker and more
pubescent than usual.

12

1513

Lippia nodiflora (L.) Mx.

12

1514

Ambrosia psilostachya DC.

12

1515

Eriochloa longifolia Vasey.

Martin County.

With the preceding.

Rather dry pine barrens near
Brunswick. Pensacola R.R. about
5 miles east of Folkston.

Moist spots near same place.

With no. 1512. Apparently indigenous.

With the preceding.

AUG

NO

12 1516 *Nama corymbosa* (Macbride) Kuntze

Wth 1517. Johnston Co.

?

12 1517 *Elionurus tripsacoides* B. & K.

(3:50

p.m.)

Moist pine-barrens about 8 miles
from Folkston and 3 from
Bull Head Bluff, Camden Co.

?

15 1518 *Solanum sisymbriifolium*?
Very prickly throughout,
except flowers and fruit.
Berries red.Waste places near Rosinshawes, 5
Brunswick, Glynn Co.
(Photo. no. 145)15 1519 *Solanum*
Resembles *S. nigrum*.

Do.

5

15 1520 *Euphorbia humistrata*?
Growing in dense patches.

Do.

5

14

AUG

15 1521 *Glaucaria trinervia* (Spreng.) Mohr

With the preceding.

5

15 1522 *Tribulus cistoides* L.

8:30

a.m.

5

15 1523 *Sporoclea filicaulis* Blume
Corolla yellow. Stems trailing.

8:35

a.m.

5

15 1524 *Sesuvium*

5

15 1525 *Heliotropium curassavicum* L.

5

15

AUG 11

15

1526

Heliotropium

With the preceding

5

15

1527

Artemisia

Do.

5

15

1528

Distichlis spicata (L.) GreeneSalt marsh on east side of
Brunswick, near end of
Gloucester St.

2 or 3

15

1529

Kallstroemia maxima (L.) J. & H.Along edge of sidewalk on G St.
near to castle St., Brunswick.

10

(11:20
a.m.)

16

1530

Argemone alba Lessib.

(Photo. no. 151)

Waste places near the beach at
Ocean Pier, St. Simon's Island,
Wynne Co.

10

16

AUG

1913

16

1531a

Chara

In water flowing from artesian
well near ruins of hotel, Ft.
Simons Island.

10

16

1531

Smilax auriculata Walt.

On low trees just back of dunes,
east shore of island.

10

16

1532

Juniperus Bonariensis

Same place as the preceding.

10

16

1533

Paronychia Baldwinii (G. & G.) Chapm.

Same place, under trees, in
dry sand.

10

16

1534

Polygonum maritimum L.

Drifting sands between dunes
and beach. (Photo. no 153)

5

17

AUG

16

1535

Chenopodium

Land - to. s. - thrust of Ocean Ave,
St. Simons Island, Glynn Co.
Perhaps introduced?

10

18

1536

Quercus

A low shrub. Like no. 1457,
but with leaves green both sides

Dry pine - barrens near the
railroad just west of Dock
Junction, Glynn Co.

15

18

1536a

Polyporus gilvus Schw.

On dead branches of fallen hickory
on bank of salt water creek
about a mile beyond Dock
Junction.

10

18

1536b

Polystictus parvulus Klotzsch

On the ground in woods near
same place.

10

18

1537

Pinus heterophylla ?
(Not *P. Elliottii* Engelm.)

The prevailing tree in dry
woods along the creek.
(Photo no. 50, 5:13 p.m.)

10

(6:15
p.m.)

18

AUG

NO.

19

1538

Rhynchospora Grayii KunthDry pine-~~various~~ on wet side of
Cumb and stand near High Point.

30

19

1539

Bejaria racemosa Vent.(11.20
a.m.)Immature capsules very glutinous.
(Photos 51, 159)In dry pine-~~various~~ near top of
scrub thicket about $\frac{1}{2}$ mile west of
High Point landing, Cumb and Island.
Very abundant.

30

19

1540

Quercus myrtifolia Willd.In dense scrub thicket
(Photo no. 150)

30

19

1541

Pieris nitida (Bartr.) B. & H.

Do.

30

19

1541a

Dicranum Bonjeani Dedt.Do., on the sandy ground,
about a mile from landing
along the car track. (Photo 161)

30

19

AUG

19

1542

Ludwigia

calyx lobes soon spread at
spreading, finally deciduous.

Style scarcely longer than the ovary.

Meadow between the road and creek,
east of the junction of the road and Lake L.

(Photo. no. 164)

19

1543

Gerardia fasciculata Ell.

Intimately associated with the preceding.

19

1544

Himbristylis

Drifting sands near the creek
(Photo. no. 53)

19

1545

Spartina juncea (Ch.) Ell.

Do.

19

1546

Juncus dichotomus Ell.

About 3 ft. tall, growing in
large tufts.

Meadow between the road and Lake L.
(Photo. no. 54)

AUG

1901

19 1547 *Cyperus Nuttallii* Eddy

With the preceding.

19 1548 *Scirpus Americanus* Pers.Outer edge of dunes
(Photo no. 165)19 1549 *Limbristylis spadicea* (L.) Vahl

Do.

19 1550 *Eleocharis*Wet sandy places on inner shore
of island just above tide, near
High Point Lighthouse.21 1551 *Spartina polystachya* (Mx.) Ell.
8-10 ft. tall
(Photo no. 56)Brackish marshes of White Oak
River near White Oak Star

21

AUG

18

Camden County.

21

1552

Cyperus

W. H. 7/7

21

1553

Lythrum lineare L.

Do.

22

1554

Ludwigia alata Ell.

Do.

22

1555

Cladium effusum (Sw.) Tor.

Do.

22

1556

Aenida canaliculata

Do.

AUG

N

22 1557 *Scirpus robustus* Pursh

With to /

22 1558 *Zizania aquatica* L.
 Culms 10 ft. tall, 1 in. thick
 at base, leaves 7 ft. long,
 3 in. wide, panicle 4-5 ft. long.

Do.

Th - no 5

23 1559 *Galactia Elliottii* Nutt.

Rather dry bushy pine woods
 about $\frac{1}{2}$ mi. north of Colesburg.

15

23 1560 *Elephantopus*

Do.

15

23 1561 *Spartina maritima*

In mud at low water mark on
 right bank of Satilla River
 near Woodbine.

23

AUG

NO

23

1562

In swampy place

In swampy place
about a mile south of Washington.

0

23

1563

Polygonum arifolium L.
Not yet in flower

Do.

23

1564

Juncus Canadensis J. Gay.

Do.

23

1565

Kosteletzkya Virginica (L.) Gay.

Do., in edge of blackish marsh.

0

25

1566

Itea Virginica L.In swamp of Little Catfish Creek, 15
about a mile south of Washington.

2 24

AUG	AUG	NO	
23	25	1566a	<i>Boletinus porus</i> (Berk.) Peck.

With the fungus, in swamp
at the same place.

2	25	1567	<i>Mikania scandens</i> (L.) Willd. Leaves somewhat fleshy, the basal lobes overlapping.
---	----	------	--

In swamp of St. Mary's River
near Ft. Laramie & Cambridge.

2	25	1568	<i>Polygonum pennsylvanicum</i>
---	----	------	---------------------------------

Do.

2	25	1569	<i>Juniperus communis</i>
---	----	------	---------------------------

Do.

2	25	1570	<i>Sagittaria lancifolia</i> L.
---	----	------	---------------------------------

Do.

25

AUG.

NO.

25 1571 *Gordonia lasianthus* L.In swamp of St. Marks River,
about $\frac{1}{2}$ mile back from river.

5

25 1572 *Rhexia flouidana* NashAround small *Typha bifida* pond
about $\frac{1}{2}$ mile south of Kingsland,
Camden Co.26 1572a *Lentinus*On rotten log in St. Marks
River swamp near Honiac,
Charlton Co.

1162

26 1573 *Tillandsia tenuifolia* L.On living trunk of *Typha latifolia*
on bank of St. Marks River
near Honiac.26 1574 *Xyris*
Spikes dark, very acute.

Low sandy places in river swamp.

26

AUG

90.

26

1575

Panicum

With the preceding.

26

1575a

*Boletus*On the ground (sandy) among dead
leaves, etc. in a swamp.

26

1576

*Xyris*Moist pine-barrens just outside
of wet swamp near Honiac.

27

1577

Quercus minima (Larg.) Small

Wet woods of oak

Rather dry pine-barrens about 1/2
a mile northwest of Pitts
Mound. Honiac

27

1577a

*Polytrichum*On dead standing trunk of *Pinus
plastris* about 1/2 mile north
of Honiac no to the

27

AUG

28

1577b

*Sphagnum*Moist s. d. bank of river
River near bridge, Clark Co.

28

1578

*Rhexia maritima*Leaves and flowers mostly gone
Roots tuberiferousMuddy flats in river swamp
near same place. (Clark Co.)

28

1579

Eupatorium

With scaly rootstocks

So.

28

1580

Eragrostis refracta (Muhl.) Scribn.Sandy bank of river.
Perhaps common.

28

1580a

Odontoschisma prostratum (Sw.) Trevis.On roots of old trees on
bank of river.

AUG. 29

29 1580b { *Archilejeunea dypeata* (Schw.) Sch.?
Trullama Donnellii ?

On smooth bark of live
 of *Naupaka* in a swamp
 of Suwannee River about a
 mile west of Fargo, Clinch Co.

29 1581 *Carex* sp.

Same place.

29 1582 *Najas rugosa* (Vitt.) Grunty

Edge of swamp, a short distance away.
 Clinch Co.

29 1582a *Sphagnum Trinitense* B. Mull.

Lying flat on the ground in
 muddy flats of Suwannee River
 swamp near Fargo, Clinch Co.
 (Same place as nos. 15798 15799)

29 1582b *Samolophyllum adnatum* (Mx) E. L. Butt.

On bark of *Taxodium* near same
 place, above high water mark.

29

AUG

29 1582 *Gullania Donnellii* Aust.
Archilejeunea conchifolia Evans

On smooth bark of small
 saplings of *Aspen* in
 swamp of *Aspen* in
 above range, Clark Co.

29 1583 *Scleria trichopoda* C. Wright.
 Sheaths broadly winged.

Most shaded places in same
 swamp.

29 1584 *Xyris neglecta* Small

Do.

29 1585 *Anastrophus paspaloides* (Max.) Nash

Do.

(See next few pages.)

Summary

{ Algae (Chara)	1
{ Fungi	6
{ Hepaticae	4
{ Musci { Sphagnum	7
{ Other mosses	4
Spermatophytes	122
Total	144

Families represented by more than one genus.

	Genera	Species
Cyperaceae	12	20
Gramineae	8	14
Compositae	5	8
Fagaceae	2	5
Leguminosae	5	5
Ericaceae	2	4
Umbelliferae	2	3
Coniferae	2	3
Amarantaceae	2	2
Zygophyllaceae	2	2

Genera represented by more than one number. (re-matching only)

Juncus	4	Holcus	2
Ludwigia	4	Helictidium	2
Ayris	4	Fimbristylis	2
Rhynchospora	3	Juncus	2
Scleria	3	Scirpus	2
Juncus	3	Rhex	2
Polygonum	3		
Spartina	3		
Pieris	3		
Sagittaria	2		
Harzania	2		
Panicum	2		
Eryngium	2		
Dichromena	2		
Hypericum	2		
Habenaria	2		
Tillandsia	2		
Eupatorium	2		

LIST OF SPECIES

SEEN IN

OKEFINOKEE SWAMP

AUGUST 6-8

NAME

*Pallavicinia Lyellii**Odontoschisma prostratum* (Sw.) Lewis*Sphagnum**Sphagnum**Sphagnum**Leucobryum glaucum* (L.) Schimp.*Sphagnum*

HABITAT

Banks of canal

On cypress knee in Jackson Bay (no. 1482 b)

Bog near Bugaboo Landing (no. 1473 a)

somewhat drier bog near Bugaboo Island (no. 1476 a)

Jackson Bay (no. 1481 a)

Jackson Bay (no. 1481 b)

Jackson Bay (no. 1482 a)

Osmunda cinnamomea L.

not common.

Osmunda spectabilis Willd.

Along canal, not common.

Lorinseria acolata (L.) Presl.

Along canal, in shaded places.

Anchistea Virginica (L.) Presl.

Banks of canal

Very abundant in sphagnum bogs on
Barbados and island, the fronds are very wet.
Jackson Bay.

Pinus Elliottii Engelm.

- In sphagnum bogs, where it is often the
only tree over considerable areas.
Abundant all over Barbados Island
Barbados Island, not common.

Pinus serotina Mx.

Taxodium imbricarium (Nutt.) Harper

Everywhere except in prairies and
on islands. Dense forests of
this species are the so-called "swamp."

Sagittaria

Sagittaria

Syntherisma sanguinalis (L.) Vahl

Panicum digitarioides Carpenter

Sulichium arundinaceum (L.) Britton

Eleocharis elongata Chapm.

Eriophorum Virginicum L.

Rhynchospora corniculata (Lam.) Gray

Along canal.

Bog near Bugaboo Landing (no. 1473)

At old commissary at inner
end of canal (introduced)

Everywhere along canal, abundant.
Chase Prairie and Jackson Bay.

Banks of canal.

Bogs between Bugaboo Landing

In canal, mostly stems

Sphagnum bog between Bugaboo Landing
and Bugaboo Island (no. 1474)

Frequent along canal

Chase Prairie, also in small prairie
3 or 4 miles from camp

Rhynchospora

Scleria trichopoda C. Wright

Serenoa seculata (Mx.) P.

Peltandra sagittaeifolia (Mx.) Morong

Peltandra Virginica (L.) Kunth

Crontium aquaticum L.

Xyris fimbriata Ell.

Xyris

Sphagnus sp. near Bugaboo Island
(no. 1477)

Chase Prairie (no. 1486)

Abundant all over Bugaboo Island,
and extending a short distance into
the surrounding sphagnum bogs.
Bog near Bugaboo Landing, about
about half way to island.

Between landing and island (rare).
Chase Prairie and Jackson Bay.

Frequent along canal.
Chase Prairie

Along canal
Jackson Bay

In small prairie about 3 miles
west of Camp Cordia (no. 1484)

Eriocaulon compressum Lam.

Eupatya flavidula (Hb.) Kunze

Gillandsia usneoides L.

Pontederia cordata L.

Smilax laurifolia L.

Gyrotheca tinctoria (Walt.) Sal.

Limodorum tuberosum L.

Chase Prairie
Jackson Bay.

Chase Prairie

On almost every tree in it

Chase Prairie

Climbing trees in timber / Chase
very common.

Along canal. }
Chase Prairie. } Not common.

Between Bugaboo Landing and
Bugaboo Island,

Quercus minima (Larg.) Small

Chenopodium

Castalia

Magnolia glauca L.

Persea pubescens (Pursh) Larg.

Sarracenia minor Walt.
Leaves and scapes 24-44 inches tall.

Sarracenia psittacina Mx.
Considerably larger than usual.

Bugaboo Island

Bugaboo Landing (introduced)

Chase Prairie

Border of Chase Prairie
Jackson Bay.

Bank of canal. Jackson Bay
Bogs between Bugaboo Landing and Chase Prairie.

Frequent along canal
Bog near Bugaboo Landing
Near Bugaboo Island (no. 1476)
Chase Prairie and Jackson Bay.

Bog near Bugaboo Island (no 1478)

Grossera intermedia Hayne

Grossera

Itea Virginica L.

Aronia arbutifolia (L.) Ell.

Galactia Elliottii Nutt.

Rhus radicans L.

Cynilla racemiflora L.

Olex cassine L.

Bog near Bugaboo Landing (no. 147.)

Chase Prairie

Border of canal. Jackson Bay.

Bogs between Bugaboo Landing and
Bugaboo Island.

Banks of canal at ...

Indian mound on Bugaboo Island,
perhaps introduced.

Border of Chase Prairie.
Jackson Bay.

Along canal.
Jackson Bay.

Along canal
Jackson Bay (no. 1482)

Ilex glabra (L.) Mill.

Ilex lucida (L.) Mill.

Acer rubrum

Vitis rotundifolia var.

(Cultivated scuppernon reverting
to the wild state)

Gordonia Lasianthus L.

Triadenum Virginicum (L.) Raf.

Decodon verticillatus (L.) Ell.

Ludwigia

A form of *L. capitata* with very
loose elongated inflorescence

Dogs between Bagaboo Landing and
Bagaboo Island, etc.

Along canal

Along canal, in wooded area

Banks of canal. Artificially introduced,
from seeds thrown out from
boats. Frequent.

Banks of canal

Dog near Bagaboo Landing
Chase Prairie. Banks of

Banks of canal

Dog near Bagaboo Landing

Bank of canal, Chase Prairie (etc.)

Myriophyllum laxum Shuttle.

Hydrocotyle umbellata L.

Nyssa biflora Walt.

Clethra

Kalmia hirsuta Walt.

Leucothoe

Pieris nitida (Pursh.) B. & Z.

Seen only in the canal. But
doubtless grows in the prairie.

Bog near Bugaboo Landing (1471)

Dense swamp near commissary,
at end of canal.

Bank of canal

Bugaboo Island.

Bogs between landing and island.

Bogs between landing and
Bugaboo Island
Jackson Bay.

Pieris phillyreaefolia (Hook.) D.D.

Xolisma
About knee-high.

Gaylussacia dumosa (Andr.) T. & G.

Gaylussacia

Vaccinium nitidum Andr.

Limnanthemum aquaticum (Walt.) Britton

Bog near Dugaboo Landing
" " " Island (no. 1475)

Near canal at edge of Chase
Prairie, climbing high on
Taxodium imbricarium (no. 1479)
Jackson Bay

Dugaboo Island.

Dugaboo Island.

Dugaboo Island.

Dugaboo Island

In canal
Chase Prairie

Euscuta compacta Juss.

On shrubs along canal.

Gerardia linifolia Nutt.

Western edge of Chase Prairie,
near camp no. 2.

Utricularia juncea Vahl.

Chase Prairie (no. 1481)

Utricularia purpurea Nutt.

Chase Prairie

Cephalanthus occidentalis L.

Border of Chase Prairie

Eupatorium capillifolium (Lam.) Small.

Luxuriant on banks of canal
(Doubtless introduced).

Chrysopsis graminifolia (Mx.) Nutt.

Indian mound on Bagaboo Island
(Perhaps introduced).

Erechtites hieracifolia (L.) Raf.

Banks of canal at commissary
and elsewhere. (Probably introduced).

I am among the 100,000.

[illegible]

Pyraceae	1
Leguminosae	1
Hamamelidaceae	1
Scrophulariaceae	1
Rubiaceae	3
Asteraceae	1
Urticaceae	1
Theaceae	1
Hypericaceae	1
Lythraceae	1
Onagraceae	1
Kaloragidaceae	1
Umbelliferae	1
Cornaceae	1
Althraceae	1
Ericaceae	5
Querciniaceae	3
Gentianaceae	1
Auscutaceae	1
Scrophulariaceae	1
Lentibulariaceae	2
Rubiaceae	1
Compositae	3
Total	81

ECONOMIC PLANTS

ART

COMMON NAMES,

An incomplete list of the
plants known and used
by the inhabitants of the
Okefinokee Swamp regions

Information obtained mostly in the
vicinity of Folkston, Camp Cornelia
and Moniac.

TECHNICAL NAME.

Pinus palustris Mill." *resinosa* Ait.*Pinus distichum* (L.) Rich. d.*Parodium imbricatum* (Vatt.) Harper.*Agave americana**Arundo stricta**Eriophorum Virginicum* L.*Stenophyllus Floridanus* Britton.*Serenia serrulata* (L.) P. & R.

COMMON NAME, AND USE, IF ANY.

used for building & fuel.

" " " " " "

Not sufficiently abundant to be of much importance.

St. White Oak a hollow hearted tree was seen in one of a small water trough.

"Cypress". An important timber tree. Considerable quantities for building taken out of Oklawaha.

"Sage". A small tree used in Altavilla.

"Rice-grass". Some of the best of the rice-grass is made into matting.

"Indian Cotton".

"Water-grass". A small plant cultivated in fields. Palmetto or sometimes "Palm-leaf".

4-4

probably also *P. virginica*
Cratichneumon aquaticum
Xyris (any species with large compact fls.)

Eriocaulon decangulare.

T. B. via *indivisa*

Polka - Canzoni

Quercus batesbaci

" cinerea

" digitata

" Virginiana

Gueltia Florida

"If you have a little to spare, I'll
 like to have it for a
 'fire' (I should say 'fire')
 and 'brots'."

'a thors.' The birds are said to
be sometimes made into bottles
by mixing them with earth.

"Black-jack."

"²⁰ Turkey Oak."

"Red Oak",

"Live Oak".

"Poor land weed."

Magnolia glauca

Passer pubescens

Cracca Virginiana

Galactia Elliottii

Polygala ramosa

Olex leassine

"A."

"B."

"C." "D." "E." "F." "G." "H." "I." "J." "K." "L." "M." "N." "O." "P." "Q." "R." "S." "T." "U." "V." "W." "X." "Y." "Z." "a." "b." "c." "d." "e." "f." "g." "h." "i." "j." "k." "l." "m." "n." "o." "p." "q." "r." "s." "t." "u." "v." "w." "x." "y." "z." "A." "B." "C." "D." "E." "F." "G." "H." "I." "J." "K." "L." "M." "N." "O." "P." "Q." "R." "S." "T." "U." "V." "W." "X." "Y." "Z." "a." "b." "c." "d." "e." "f." "g." "h." "i." "j." "k." "l." "m." "n." "o." "p." "q." "r." "s." "t." "u." "v." "w." "x." "y." "z."

"A." "B." "C." "D." "E." "F." "G." "H." "I." "J." "K." "L." "M." "N." "O." "P." "Q." "R." "S." "T." "U." "V." "W." "X." "Y." "Z." "a." "b." "c." "d." "e." "f." "g." "h." "i." "j." "k." "l." "m." "n." "o." "p." "q." "r." "s." "t." "u." "v." "w." "x." "y." "z."

and to yield a yellow eye.

"Zassena" These are considered
("Henderson Wood") distinct species,
but apparently the only character
which the natives use to separate
the "Henderson Wood" is the
size, and the spots on the
back, the color of the
feathers, and the different sizes
simply correspond to different ages.

Sida rhombifolia

Gordonia lasianthus

Eriadenum virginicum

"*St. John's wort*". The dead stems are
used to retain water, (like
the information is altered
at the cost in the plant).

"*St. John's wort*". This is frequently used
in the habitat with the other
two "worts", being in
the same family as the
other two.

"*St. John's wort*". (Known as in
Okefinokee Swamp). The origin of
this name is a mystery. It
seems scarcely credible that
these uneducated people should
know that this plant belongs
to the same family as
"*St. John's wort*", "*St. Peter's wort*",
etc.

to illustrate the system
of the ...
according to the ...

Trees

Evergreen (Conifers)

Deciduous

Shrubs

Evergreen

Pinus Elliottii" *resinosa**Juniperus imbricaria**Thuja occidentalis**Yucca filifera**Quercus minima**Rapanea glauca**Persea pubescens**Leucaena*" *glabra*" *indica**Albizia leucodermis**Albizia leucodermis**Albizia leucodermis*" *philippensis**Polium**Gaylussacia dumosa*" *Vaccinium nitidum*

Shrubs (continued)
Deciduous

Palms, acaulescent

Woody vines, climbing
Rootlets

Tendrils

Creeping under bark of *Taxodium*

at
A. ...
...
...
...
...
...
...

Scorpaenaria

Rhus radicans

Smilax laurifolia

Pieris phillyreaefolia

Herbs

Parasitic

Epiphyt.

Free floating

Attached to the water
aquatic
float. floating leaves


Etc.

as the composite

as the composite

as the composite

Castalia
Linnanthemum aquaticum



II
LIST OF PLANTS

OBSERVED IN AND NEAR
OKEFINOKEE SWAMP,
GEORGIA

IN THE COUNTIES OF
WARE, PIERCE, WAYNE, GLYNN,
CAMDEN, CHARLTON, CLINCH,
ECHOLS AND LOWNDES
IN AUGUST, 1902,
BY
ROLAND M. HARPER.

ALGAE

Chara

FUNGI.

Polystictus parvulus Klotzsch.

Polyporus ...

Polyporus gilvus Schw.

Boletus ...

Boletinus porosus (Berk.) Peck

Lentinus

In pool of fresh water around
artesian well, St. Simon's I. (1530a)

On the ground in dry wood near
Dock Junction, Glynn Co. (1536b)

On dead *Pinus palustris* near
Moniac, Charlton Co. (1577a)

On dead *Glicoria* near Dock
Junction, Glynn Co. (1536a)

In St. Mary's River swamp near
Moniac, Charlton Co. (1575a)

On rotten logs in swamp of Little
Creek, Camden Co. (1566a)

On rotten logs in St. Mary's River
swamp near Moniac, Charlton
Co. (1572a)

HEPATILAE

Pallavicinia Lyellii (Hook.) S. F. Gray

Odontoschisma prostratum (Sw.) Trevis

Archileporeia dyceata (Schw.) Hiffn.

" *conchifolia* Evans

Grullania Donnellii Aust.

MUSCI.

Sphagnum Ludovicianum (R. & G.) Warnst.

" *Portoricense* Blampe

" *tenerum* (Aust.) Warnst.

On the ground in swamp of Little
Calfish Creek,

Jackson Bay, Okefinokee Swamp. (1482b)
Bank of Suwannee River. (1580a)

Swamp of Suwannee River, Echols Co. (1580b)

Swamp of Suwannee River,
Blinch Co. (1582c, in part)

Intimately associated with the
two preceding (1580b, 1582c).

Near Glenmore, Ware Co. (1470a)
Okefinokee Swamp, near Bugaboo Q. (1476a)

Okefinokee Swamp, Jackson Bay (1476a)

Bog east of Holston (1508a)

Sphagnum Trinitense B. Hill.

"

"

Dicranum Bonjeani De Not.

Leucobryum glaucum (L.) Schimp.

Sematophyllum adnatum (Mx.) C. G. Britton

Myrica

Swannee River swamp near
Fargo, Clinch Co. (1582a)

Dog near Bugaboo Landing, Okefinokee
Swamp (1473a)

Sandy bank of Swannee River
near Fargo, Clinch Co. (1577b)

On the ground in scrub thickets,
Cumberland Island, (1541a)

Okefinokee Swamp, Jackson Bay (1481b)

On bank of *Taxodium*, Swannee
River swamp near Fargo (1582b)

Okefinokee Swamp, Jackson Bay (1482a)

PTERIDOPHYTA

OSMUNDACEAE.

Osmunda cinnamomea L.*Osmunda spectabilis* Willd.

POLYPODIACEAE

Lorinseria areolata (L.) Presl.
(*Woodwardia areolata* (L.) Moore)*Anchistea Virginica* (L.) Presl.
(*Woodwardia Virginica* (L.) J. E. Smith)*Athyrium Filix-femina* (L.) Roth{ In swamps. Camden: Near
Woodbine; along Little Batfish Cr. & St. Mary's River.
Charlton: Near Folkston; Okefinokee Swamp.{ Camden: - Swamps of Satilla and
St. Mary's Rivers and Big Batfish Creek.
Charlton: Near Folkston; Okefinokee Swamp.{ Shaded swamps. Camden: West
of Woodbine; Little Batfish Creek.
Charlton: Near Folkston; Okefinokee Swamp.{ Wet pine-barrens and non-
alluvial swamps, common throughout,
except immediately along the coast.
Ware, Pierce, Glynn, Camden,
Charlton, Blinch, Echols, Lowndes.
Particularly abundant in
portions of Okefinokee Swamp.{ Camden: Wet woods around spring
south of Kingsland.
Charlton: Bluff at Traders Hill.

Pteridium aquilinum

Polypodium polypodioides (L.) Hitchcock

LYCOPODIACEAE.

Lycopodium alopecuroides L.

" *Chapmani* Underw.
(*L. alopecuroides adpressum* Chapm.)

Lycopodium Carolinianum L.

" *pinnatum* (Chapm.) L. & C.

Dry pine-barrens and sand-hills.
Pierce: along Saluda. Glynn:
Near Dock Junction. Camden: Woodbine.
Charlton: Folkston; Ridge near Camp Cornelia.
Camden: Cumberland Island; St. Marys I. swamp.
Charlton: swamps west of Folkston.

Moist pine-barrens.
Ware, Camden, Charlton, Bluff.

Charlton: Moist sandy place
just south of Folkston.

Charlton:— Moist pine-barrens
between Folkston and Camp
Cornelia; Bog east of Folkston.

Ware: Moist pine-barrens
between Waycross and Saluda R.;
Rather dry pine-barrens just
west of Waycross.

GYMNOSPERMAE

Pinus Elliottii Engelm.

Common throughout in low pine-barrens. Seen in every county. Extends eastward to Cumberland Island, nearly to Brunswick, and perhaps on Jekyll Island. Forms exclusive forests in sphagnum bogs of Okefinokee Swamp, and occurs similarly in sphagnum swamp of St. Charles River south of Waycross.

Pinus palustris Mill.

Common throughout in dry pine-barrens. Seen in every county, but not in Okefinokee Swamp or in any maritime association. Extends eastward to Cumberland Island (High Point) and to within 7 or 8 miles of Brunswick.

Pinus serotina Mx.

Moist pine-barrens and sandy swamps. Seen in every county except Echols, but nowhere common, except near Glenmore. Extends east to about Dock Junction. Occurs sparingly on Bugabo Island in Okefinokee Swamp.

Pinus

Pinus

Taxodium distichum (L.) Richard

Camden: White Oak.

Charlton: Bluff at Traders Hill.

Clinch: Along Suwannee River and Suwannee Creek near Fargo.

Glynn: Cypress woods near Dock Jc. (1537); also between Cartier and Thalmann.

Camden: Along Satilla River near Woodbine; and a small creek north of Colesburg.

Charlton: Near Spanish Creek; along river at Traders Hill.

Glynn: Swamps along railroad from Thalmann southward.

Camden: Along Satilla River at Bull Head Bluff and Woodbine; Big Catfish Creek; St. Mary's River; Swamps near Glenora and White Oak.

Charlton: Along St. Mary's River near Folkston, Traders Hill and Noriac; swamps between Folkston and Bull Head Bluff.

Clinch: On and along Suwannee River near Fargo.

Taxodium imbricarium (Vitt.) Blayden

Juniperus ?

ANGIOSPERMAE

MONOCOTYLEDONES

TYPHACEAE

Typha latifolia L.

NAIADACEAE

Potamogeton diversifolius Raf.

"

ALISMACEAE

Sagittaria lancifolia L.

Common throughout in wet pine-
barrens, ponds, etc. Seen in
every county. Particularly abundant
in Chefinokee Swamp. In
Satilla River as far down as
Waycross. Extends east to just
beyond Dock Junction.

Olynn: Among the dunes on Ft.
Simons Island (1532)

Camden: St. Mary's River swamp (1569)

{ Fresh marshes.

Olynn: Between Dock Jc. and Bartier

Camden: Swamps of White Oak and Satilla Rivers

Lowndes: Ten miles west of Valdosta

Ware: Cypress pond about 3 miles
southwest of Waycross.

Camden: Wet woods near cool spring
south of Kingsland

{ Camden: Swamps and marshes of
White Oak, Satilla & St. Mary's (1570) Rivers

Charlton: Along river at Traders
Bluff (6 ft. tall.)

Sagittaria latifolia Willd.

Sagittaria Mohrii J. G. Smith

Sagittaria

HYDROCHARIDACEAE

Limnobium Spongia Richard.

GRAMINEAE

Manisuris rugosa (Nutt.) Kunth.

Eleocharis tupsacoides H. & A.

Andropogon tener Kunth.

Anastrophus compressus (Sw.) Schlecht.

" *paspaloides* (Mx.) Nash

Camden: Edge of Satilla River marshes
at Woodbine.

Chefinokee Swamp. Valdosta.

{ Ware: Wet places in railroad right-
of-way between Waycross and
Chefinokee Swamp; Log 12. Bugaboo
Landing. (1473)

Camden: Pools near railroad south
of Colesburg and Kingsland.
Perhaps not indigenous.

{ Charlton: Moist pine barrens about 5
miles east and west of Charlton;
Between Butler and Moniac.

Echols: Edge of swamp of Suwannee
Creek (1582); Near Howell.

{ Camden: Moist pine-barrens west of
Bull Head Bluff (1517)

Lowndes: Pine-barrens east of Valdosta.

Ware: Along railroads near Waycross. Introduced

Ware: Moist pine-barrens between Waycross & river.

Charlton: River swamp near Moniac

Chilch: " " " Fargo (1585)

Lowndes: Valdosta.

Paspalum distichum L.

" *Boscianum* Flugge

" *setaceum* Mx.

Paspalum

Syntherisma sanguinale (L.) Nash

Panicum digitarioides Carpenter

Panicum

" *barbulatum* Mx.

" *gibbum* Ell.

" *scabriusculum* Ell.

Alyssa: Ground artesian well, St. Simon's I.
Charlton: ditch along inland south of F. Astor.
Camden: Cornfield west of Bull Head Bluff

Lowndes: Low grounds, Valdosta.

Charlton: Shallow cypress pond near
 Camp Cornelia (1486)

A common weed, found even in
 the middle of Okefinokee Swamp,
 along the canal.

Brunswick, Traders Hill, Valdosta, etc.

Ware: Pine-baren ponds near Glenmore
Charlton: West of Butler.

All along canal in Okefinokee Swamp,
 also in prairies and bays.

Lowndes: Cypress ponds, Valdosta, and eastward
Alyssa: Dry pine-barens near Stock Is.

Camden: Damp woods, White Oak.

Camden: Along Little Catfish Creek.

Swamps. *Ware*: Near Ruskin

Charlton: First *Taxodium distichum*
 swamp east of Collier.

Panicum stenodes Griseb.

Panicum virgatum L.

Panicum

Panicum

Panicum

Panicum

Cenchrus

Echinochloa colona (L.) Link.

Eriochloa longifolia Vasey.

Oplismenus setarius (Lam.) R. & S.

Zizania aquatica L.

{ Ware: Cypress ponds near Waycross and Abbeville

{ Charlton: " " north of Camp Cornelia
Spanish Creek flats.

Dry pine-barrens, Cumberland Island.

{ Camden: Wet woods near spring
south of Kingsland.

Camden: Satilla River swamp near Woodbine

{ Charlton: Shallow cypress pond
south of Camp Cornelia (1485)

Charlton: River swamp near Moniac (1575)

{ Sweet: Waycross, Brunswick
and Camp Cornelia

Among wharves, Brunswick.

{ Camden: Moist pine-barrens west
of Bull Head Bluff.

Charlton: A few miles farther west (1515)

{ Glynn: Dry thicket back of dunes,
St. Simon's Island.

{ Camden: Rich woods near cool
spring south of Kingsland.

{ Camden: Brackish marshes of White
Oak River (1558) and Big Batfish Cr.

Aristida palustris (Chapm.) Vasey

Aristida spiciformis Ell.

Aristida stricta Mx.

Sporobolus Indicus (L.) R. Br.

" *gracilis* (Trin.) Merrill

Cappirola Dactylon (L.) Kunth

Spartina juncea Mx. Ell.

Spartina maritima

{ Shallow pine-barren ponds.

Ware: Near Beach, in northern part of county.
Charlton (?): Northwest of Elliott's Pond.

{ Flat or rather dry pine-barrens.

Ware, Pierce, Camden, Charlton,
Blinch, Echols, Lowndes.

{ Dry pine-barrens, wherever these occur.

Pierce: Vicinity of St. Mudge.

Glynn: " " Dock Junction

Camden: South of Kingsland

Charlton: On ridge near Camp Cornelia.

Streets of Brunswick, etc.

{ Dry or rather dry pine-barrens.

Camden: Cumberland Island

Blinch: Near Fargo.

{ A weed, sometimes escaped from cultivation.

Glynn: Brunswick; St. Simon's Island

Blinch: Fargo. Echols: Mayday.

{ Drifting sands near the beach,
Cumberland Island. (1545)

Glynn: Salt marshes on St. Simon's
Island, and near Brunswick and
Dock Junction.

Camden: In Satilla River at Woodbine;
a remarkably slender form (1561)

Spartina polystachya (Mx.) Ell.

Campulosus aromaticus (Walt.) Trin

Eleusine Indica (L.) Gaertn.

Dactyloctenium aegyptium (L.) Moench

Sieglingia ambigua (Ell.) Kunze.

Eragrostis simplex Scribn.

Eragrostis refracta (Muhl.) Scribn.

Uniola paniculata L.

" *laxa* (L.) B. & P.

Distichlis spicata (L.) Greene

{ Camden: Brackish marshes of
White Oak (1551) and Hatilla
and Big Batfish Creek.

{ Moist or rather dry pine-land.
Ware, Glynn, Camden, Charlton,
Clinch, Lowndes. In Charlton
occurs also in the Rhexia glabella flats.
Not seen farther east than S. A. L.

{ A weed. White Oak and Folkston,
around dwellings.

{ A weed of sandy cultivated fields.
Traders Bluff, Valdosta, etc.

{ Charlton: Rhexia glabella flats of
St. Mary's River and Spanish Creek.
Between Folkston and Bull Head Bluff.

{ Along railroads, especially between
the rails. Common. Noted near
Waycross, Kingsland and Moultrie.
Sandy bank of Suwannee River (1580)

{ On dunes. South end of Jekyll Island
and north end of Cumberland.

{ Camden: Swamp of Little Batfish Creek

{ Charlton: " " St. Mary's R. near Moultrie

{ Echols: " " Suwannee Creek.

{ Glynn: Salt marshes, Brunswick (1528)

{ Camden: Inner shore of Cumberland.

CYPERACEAE

Lipocarpus maculata (Mx.) Torr.*Cyperus compressus* L.

"

" *filiculmis* Vahl." *flavescens* L." *haspan* L." *iria* L." *praelongatus* Steud." *Nuttallii* Eddy" *rotundus* L." *squarrosus* L." *virens* Mx.

Low grounds, perhaps introduced
 Charleston: Between Folkston and Kingsland.
 Lowndes: South of Valdosta.
 A weed. White Oak, Moniac, Fargo.

Meadow near beach, Cumberland Island.

Dry pine barrens and sand dunes.
 Glynn: Near Dock Junction.
 Charleston: Near Folkston, and on
 ridge near Camp Cornelia.

Around artesian well, St. Simon's Island.

Bamden: Brackish marshes of White Oak R.
 Charleston: Moist sand south of Folkston.
 Echols: Near Suwannee Creek.
 Lowndes: Valdosta, and southward.

In cultivated ground, White Oak (rare)

Brackish marshes of White Oak River (1552)

Meadow near beach, Cumberland I. (1552)

Streets of Brunswick

A weed. Moniac and Fargo.

Brackish marshes of White Oak River.

Scirpus arundinaceus (L.) Britton

Eleocharis baldwinii (Torr.) Chapm.

Eleocharis elongata Chapm.

" *interstincta* (Vahl) R. & S.

" *melanocarpa* (Baldw.) Torr.

" *tuberculosa* (Mx.) R. & S.

Eleocharis

Dichromena colorata (L.) Hitchcock.

"

A slender form, perhaps distinct sp.

Charlton: Moist sandy places south of
Folkston, between Norcross and S. Ma.
and northwest of Ellicott's Bend.
Also in Okefinoche Swamp.

Rather dry pine-barrens, especially
abundant along paths and unfrequented roads.
Ware, Glynn, Camden, Lowndes,
Chatham, Chocoma. Extends east to
within 10 miles of Brunswick.

In canal, Okefinoche Swamp.

{ Camden: Shallow pond south of
Lowndes: " " " " Valdosta
Charlton: Sand-hill pond southwest of Folkston.
Moist pine-barrens and ditches.
Ware: Between Waycross and Ruskin.
Camden: " " " " Seals.
Charlton: Along R.R. south of Folkston.

{ Camden: Inner shore of Cumberland and S.
(1550); Brackish marshes of W. to Oak R.
Camden: Meadow on Cumberland; White Oak marshes.
Charlton: Along River at S. Hill (1500)
" Flats near Spanish Creek (1505)

Gichromena latifolia Baldw.

Stenophyllus ciliatifolius (Ell.) Britton ?

" *Floridanus* Britton

" *stenophyllus* (Ell.) Britton

Moist and usually flat pine-barrens.

Ware: frequent.

Camden: Between Kingsland & St. Mary's R.

Charlton: Near glabella flats, etc.

Glynn: Dry pine-barrens between Brunswick and Dock Junction.

Strictly ruderal, but always in dry sand (Columbia formation).

Ware: Near Ruskin.

Charlton: Folkston, Traders Hill, Moniac, etc.

Blinch: Fargo.

Echols: Howell; and abundant in cornfield about 2 miles farther west.

Townsend: Near Blanton; Valdosta.

Not observed in the maritime counties.

Habitat similar to the preceding

Glynn: Brunswick, and along railroads 3-4 and 7-10 miles west.

Camden: Between rails, on track between Woodbine and Colesburg.

Charlton: Vicinity of Camp Cornelia, Traders Hill and Moniac.

Blinch: Fargo and vicinity.

Always appears as if introduced.

Limbristylis autumnalis (L.) R. & S.

" *puberula* (Mx.) Vahl.

" *spadicea* (L.) Vahl.

"

Scirpus Americanus Pers.

" *Eriophorum* Mx.

" *lacustris* L.

Low grounds, ponds, ditches, etc.

Camden: Exiccated ponds south of Kingsland.

Charlton: West of Spanish Creek: Along river at Traders Hills.

Lowndes: South of Valdosta.

Dry or rather dry pine-barrens.

Ware: Just southwest of Waycross.

Charlton: Folkston, and westward to ridge; Near Moniac.

Glynn: Borders of small salt marsh just west of Brunswick.

Camden: Outer edge of dunes, Cumberland (1549)

Camden: Drifting sands near beach, Cumberland Island (1544)

Outer edge of dunes, Cumberland I. (1548)

Ditches and swamps, especially in clearings.

Ware: Near Glenmore.

Glynn: South of S.B.T. Cussing (indigenous)

Camden: Satilla River swamp near Woodbine.

Charlton: Abundant in great ditch at Camp Cornelia; *Eleocharis distichum* swamp northeast of Folkston.

Lowndes: East of Blanton; South of Valdosta.

On Satilla River at Woodbine, growing in edge of water and bathed by the tides; very tall. (Photo. 59)

Scirpus robustus Pursh

Eriophorum Virginicum L.

Guirena breviseta Coville.

"

" *scirpoides* Mx.

Rhynchospora axillaris (Lam.) Britton

"

Baldwinii Gray.

"

caduca Ell.

"

Chapmani, M. & Curtis.

Brackish marshes of White Oak River (1557)

Sphagnum bogs, Chequamegon Swamp (1474)

Moist places along railroad south of Valdosta.

Moist pine-barrens. Ware: Between
Waycross and Satilla River.

Lowndes: Valdosta

Ditches along railroads near Folkston & Valdosta.
Rather dry pine-barrens north and
east of Camp Cornelia.

Moist pine-barrens and pond-margins.

Ware: Between Waycross and Rushin.

Camden: Woodbine

Charlton: West of Spanish Creek;
Northwest of Olicott's Mound.

Echols: Near Suwannee Creek.

Lowndes: Valdosta.

Ware: Moist pine-barrens between
Waycross and Satilla River.

Charlton: *Rhexia glabella* flats along
St. Mary's River south of Folkston.

Clinch: Low woods near Suwannee Cr.

Ware: Moist pine-barrens near Rushin.

Rhynchospora ciliaris (Mx.) Mohr

" *corniculata* (Lam.) Gray.

" *distans* (Mx.) Vahl.

" *fascicularis* (Mx.) Vahl.

" *filifolia* Torr.

" *glomerata paniculata* (Gray) Chapm.

" *Grayii* Kunth

" *inexpansa* (Mx.) Vahl.

Moist or rather dry pine-barrens.
Ware: Between Waycross and Ruskin.
Charlton: Near Folkston, and westward

Wet pine-barrens, ponds and swamps.
Ware: North edge of county. Very abundant
between Waycross and Glenmore.

Southeast of Waycross.

Charlton: *Taxodium distichum* swamp
northeast of Folkston; Vicinity of
Ellicott's Mound; Okefinokee Swamp.

Blinch: Near Fargo, and westward
along railroad.

Charlton: Rather dry/sandy road.
Camp Cornelia (1487)

Blinch: Rather dry pine-barrens near Fargo.

Charlton: Okefinokee Swamp (1477)

Camden: Sphagnum swamp of St. Mary's R.

Charlton: Cypress pond south of Camp Cornelia

Lowndes: Low grounds south of Valdosta.

Pierce: Sand-hills of Satilla River

Camden: Cumberland Island (1538)

Glynn: Moist places, near Cartier
and Thalmann.

Rhynchospora

" *semiplumosa* Hay

Cladium effusum Sw. Torr.

Scleria Baldwinii Torr.

" *gracilis* Ell.

" *hirtella* Sw.

Camden: Swamps of Satilla and
St. Mary's Rivers and Little Catfish Cr.

Charlton: *Rhexia glabella* flats along
Spanish Cr.

Brackish marshes and moist pine-barrens
(in the latter, usually sterile)

Glynn: Near Dock Junction

Camden: Pine-barrens west of Bull Head Bluff;

Between Woodbine and Colesburg;

Marshes of the three rivers (luxuriant)

Charlton: Various points east of Holston.

In and around cypress ponds.

Ware: Between Waycross and Ruskin

Charlton: South of Camp Cornelia

Northwest and east of Ellicott's.

Not seen east of Apalachicola Ridge.

Charlton: *Rhexia glabella* flats of

St. Mary's River and Spanish Cr.

Cypress pond south of Camp Cornelia.

Charlton: *Rhexia glabella* flats of

St. Mary's River and Spanish Creek;

4 or 5 station between " "

and Camp Cornelia (1492), in

moist pine-barrens.

Scleria trichopoda B. Wright.

"

"

Carex

"

PALMAE

Sabal glabra (Mill.) Sarg.
(*S. Adansonii* Swens.)

Sabal Palmetto

Chefinokee Swamp: Chase Prairie (1480)

Glinch: Suwannee River swamp (1583)

Lowndes: Moist pine-barrens, Valdosta.

Not seen east of Chefinokee Ridge.

Lowndes: Moist pine-barrens, Valdosta.

Charlton: Sand-hills southwest of Folkston.

Camden: Swamp of Little Catfish Creek.

Moist pine-barrens and swamps.

Ware: Between Waycross and Rushin

Charlton: Vicinity of Folkston, especially about 5 miles northeast; Near Monice.

Echols: Swamp of Suwannee Creek (1581)

Glynn: Woods west of Dock Junction

Camden: " between Woodbine and Satilla R.

Swamp " " " Bolshung.

St. Mary's River swamp (abundant)

Charlton: " " " bank at Traders Hill

Glynn: Borders of salt marshes near Brunswick; Marshes and pine-barrens along B. & O. R. R. from 13th mile post to Thalmann; Abundant (tall and slender) in pine-barrens around Thalmann and southward to S. B. I. Crossing.

Serenoa serrulata (Mx.)
(*Corypha repens* Bartr.)

ARACEAE

Peltandra sagittaeifolia (Mx.) Morong.

" *Virginica* (L.) Kunth.

Orontium aquaticum L.

MAYACACEAE

Mayaca dubletti Mx.

XYRIDACEAE

Xyris Baldwiniana R. & S.

Common throughout, from the dunes of the sea-coast westward, from swamps to pine-barrens & the driest sand-hills. Seen in every county. Completely covers Bugaboo Island in Okefinokee Swamp. But absent from many places which seem perfectly adapted to it. Very rarely fruiting.

{ Sphagnum bogs near Bugaboo Landing,
Okefinokee Swamp.

{ Camden: Satilla River swamp near Woodbine.
Charlton: Bogs, prairies and bays of
Okefinokee Swamp.

{ Charlton: Swamp about 3 m. E. of Folkston.
Common in Okefinokee Swamp.

Charlton: Sphagnum bogs near Folkston

Charlton: Sandy bogs near Folkston.

Xyris brevifolia Mx.

" *elliottii* Chapm.

" *fimbriata* Ell.

" *neglecta* Small.

" *smalliana* Nash.

" *torta*, J. E. Smith.

"

Rather dry pine-barrens, often around ponds.

Ware: Between Waycross and Ruskin.

Charlton: Several stations near Camp Cornelia.

Blind: Southwest of Fargo.

Not seen east of Okefinokee Ridges.

Habitat similar to the preceding.

Ware: Between Waycross and Ruskin.

Camden: Small pond south of Big Catfish Cr.

Charlton: Near Camp Cornelia.

Pine-barren ponds, etc.

Camden: South of Big Catfish Creek.

Charlton: Northwest of Ellicott's Mound,
and in Okefinokee Swamp.

Lowndes: Near Stanton and Valdosta.

Blind: Suwannee River swamp (1584)

Okefinokee Swamp (1484)

Rather dry pine-barrens.

Ware: Near Waycross.

Camden: Woodbine; south of Kingsland.

Charlton: East of Folkston, and on
ridge near Camp Cornelia.

Lowndes: South of Valdosta.

Charlton: Moist pine-barrens near Moultrie
(1510)

Xyris

ERIOCAULACEAE

Eriocaulon compressum Lam.

" *decangulare* L.

Lachnocaulon anceps (Walt.) Morong

" *longicaule* (Lam.)

Dupatya flavidula (Mx.) Kuntze

Charlton: River swamp at Moniac (1574).
Clinch: " " " Fargo

Charlton: Cypress pond south of Camp Cornelia;
 Prairies and bays, Okefinokee Swamp.

Moist pine-barrens, etc.; common.
 Seen in every county except Pierce
 (and doubtless occurs - there). Not
 seen farther east than S. of L. Ry.,
 or in Okefinokee Swamp

Moist or rather dry pine-barrens.

Ware: Between Waycross and Ruskin.

Camden: Woodbine.

Charlton: Common.

Charlton: Rather dry pine-barrens
 north of Camp Cornelia (1491)

Moist or rather dry pine-barrens, etc.

Ware: Near Glenmore.

Charlton: Okefinokee Swamp, and elsewhere,
 (very rarely east of the ridge)

Clinch: Southwest of Fargo.

Lowndes: South of Valdosta

BROMELIACEAE

Tillandsia tenuifolia L." *usneoides* L.

PONTEDERIACEAE

Pontederia cordata L.

JUNCACEAE

Juncus" *canadensis* J. Gay." *dichotomus* Ell." *effusus* L.

{ Charlton: Swamp east of Holkston (1579)
 Along river at Honiac (1573)
 Quite common, usually near water.
Ware, Glynn, Camden, Charlton,
Clinch, Ekols, Lowndes. Very common
 in Okefinokee Swamp.

Common in pine-barren ponds,
 ditches, and brackish marshes.
 In marshes larger (6 ft. tall on
 Satilla River at Woodbine), and
 with broader and thinner leaves.
 Common in Okefinokee Swamp.
 Seen in every county except Pierce.

{ Pierce: Ordinary sand-hills of Satilla R.
Charlton: Sand-hills near Holkston (1495)
 Pine-barrens northwest of Ellicott's mound

Satilla River swamp near Woodbine (1565)

Meadow near beach, Cumberland I. (1546)

Ware: Along Satilla River at Wattertown;
 " railroad near Ruskin
Charlton: " " " Holkston.

Juncus megacephalus, N. & Kurtz.

" *polgcephalus* Mx.

" *repens* Mx.

" *Roemerianus* Scheele.

" *trigonocarpus* Steud.

" *rostratus*

Heads - near beach, Cumberland Island.

Moist pine - barrens, pools and ditches.

Ware: Between Ruskin and Glenmore.

Charlton: About 6 miles northeast of Ft. Mifflin.

Clinch: Near Shelma.

Lowndes: South of Valdosta.

Habitat similar to the preceding.

Ware: Between Ruskin and Glenmore.

Camden: Pond south of Kingsland.

Charlton: Near Racepond, and west of Spanish Creek.

Echols: Near Mayday.

Lowndes: " Blanton.

Glynn: Common in salt marshes around Brunswick. In salt creek beyond Dock Junction (photo. no. 50).

Covers several acres near Bartow.

Camden: Inner shore, Cumberland I.

Marshes of White Oak River.

Wet pine - barrens near Racepond.

Pierce: Moist depressions in sand-hills of Satilla R.

Glynn: Around small pond near Dock Jc.

Charlton: Vicinity of Colleton River.
Camp Cornelia, several stations

LILIACEAE

Tofieldia racemosa (Walt.) D. & P.*Melanthium dispersum* Small.*Lilium batesbaci* Walt.*Yucca aloifolia* L.*Aletris aurea* Walt." *farinosa* L.

SMILACEAE

Smilax auriculata Walt." *Bona-Nox* L." *humila* Walt.Ware: Moist pine-barrens near Waycross.
Charlton. Sphagnum bog east of Holkston.

" " " " " (1506)

{ Moist pine-barrens, common, but
never more than a dozen specimens visible
at once. Ware, Pierce, Banden,
Charlton, Clinch, Lowndes.
Among the dunes, St. Simon's Island.

Rather dry pine-barrens, Waycross.

" " " " N.W. of Ellicott's Mound.

Climbing low trees just back of
dunes, St. Simon's Island. (1531).

Mixed with the preceding.

Ordinary sand-hills of Satilla River,
Pierce Co.

Smilax laurifolia L.

HAEMODORACEAE

Erythraea tinctoria (Walt.) Sal.

Lophiola aurea Ker.

IRIDACEAE

Iris versicolor L.

CANNACEAE

Canna flaccida Roscoe.

BURMANNIACEAE

Burmannia capitata (Walt.) Mart.

Common in wooded non-alluvial swamps, especially Okefinokee Swamp.

Camden: Swamp of St. Mary's River.
Charlton, Clinch, Lowndes.

Wet pine-barrens, ponds, and non-alluvial swamps.

Ware: Near Waycross and Rushin.

Camden: St. Mary's River swamp.

Charlton: Between Butler and Moniac.
Near Elliott's Mound; Okefinokee Swamp.

Ware: Moist pine-barrens near Rushin.

Charlton: Wet pine-barrens between Spanish Creek and Okefinokee Ridge.

{ In *Taxodium distichum* swamps (no Lafayette).
Camden: Between Waverly and White Oak.
Charlton: About 6 miles S.E. of Folkston.

Charlton: Moist ditch along railroad south of Folkston; Sphagnum bog about 2 m. east of Folkston.

Xabenaria blephariglossa (Willd.) Torr.

" *ciliaris* (L.) R. Br.

" *integra* (Nutt.) Spreng.

Limodorum tuberosum L.

{ Moist pine-barrens and sphagnum
bogs, not rare. Pierre, Camden,
Blairstown (1508) Lowndes.

{ Habitat similar to the preceding.
Glynn: 6-7 and 9-10 miles west of
Brunswick on B. & O. R.R.
Camden: Near White Oak; South of
Big Catfish Creek.
Lowndes: South of Valdosta.

{ Habitat similar to the two preceding.
Ware: Between Murrays and Beach.
" Waltham " Waycross.
Charlton: East (1507) and southwest
of Golleston.

Charlton: Sphagnum bogs.
About 2 miles east of Golleston,
also in Okefenokee Swamp.

DICOTYLEDONES

SAURURACEAE.

Saururus cernuus L.

JUGLANDACEAE

Alcorno

MYRICACEAE

Myrica carolinensis Mill." *erifera* L.

{ Pine-barren ponds and swamps.
Camden: Just south of Bolesburg (rare)
 Swamp of Little Catfish Creek.
Charlton: Near Holkston; along
 river at Traders Hill.

Glynn: Dry woods on bank of salt
 creek just west of Lock Junction.

{ Wet pine-barrens, especially about
 edges of non-alluvial swamps.
Charlton: South and east of Holkston.
Camden: Near Little Catfish Creek.

Habitat very variable.

Glynn: Dunes of St. Simon's; Woods near
 Brunswick, and near Cartier.

Camden: Scrub thicket, Cumberland I.
 Woods and swamps all along
 the railroad.

Charlton: In swamps with either species
 of *Taxodium*; Along St. Mary's River
 at Traders Hill (20 ft. tall)

Blinch: Low woods near
 Suwanoochee Creek, west of Fargo.

31

Myrica humilis (Mx.) Small.

BETULACEAE

Carpinus Caroliniana Walt.

Alnus rugosa (DuRoi) Koch.

FAGACEAE

Castanea nana Ell.

Castanea

Castanea

Usually in dry pine-barrens.
Glynn: Between Brunswick & Dock Jr.
Camden: South of Kingsland.
Charlton: Near Folkston & Camp Cornelia.
Pierce: Sand-hills of Satilla River;
 Vicinity of Ft. Mudge.

Wooded bluff at Traders Hill.

Charlton: Along streams at several points within a few miles of Folkston.

Dry pine-barrens.
Ware: Between Elsie and Upchurch.
Pierce: " Schlatterville & Hoboken.
Glynn: Vicinity of Dock Junction.
Camden: Colesburg.
Charlton: Near Racepond and Folkston.

Pierce: Driest sand-hills of Satilla River (1467).
Camden: Sandy woods near Satilla River at Woodbine.

Wooded bluff at Traders Hill.

Quercus alba L.

" *batesbaei* Mx.

" *brevifolia* (Lam.) Sarg.
(*Q. cinerea* Mx.)

Quercus digitata (Marsh.) Sudw.

Camden Woods near Little R. at Woodline
Lowndes: south of Camden

Sand-hills and dry pine-barrens.
Ware: Sand-hills of Satilla River
near Walbertown. Also east of Waycross.

Pierce: Ordinary sand-hills of Satilla R.

Lynn: Dry pine-barrens around Dock Jr.

Camden: Cumberland Island.
South of Kingsland.

Charlton: Vicinity of Racepond,
Folkston and Butler, common.

Echols: Both sides of Allapaha
River near Mayday.

Habitat similar to the preceding,
with which it is usually associated.

Ware: Sand-hills of Satilla R. ^{near Walbertown}

Pierce: " " " " "

Also near Schlatterville.

Camden: South of Kingsland

Charlton: Folkston and vicinity.

Echols: Along Allapaha River
near Mayday.

Charlton: Woods along streams west
of Folkston, and near Camden
Co. line northeast of there.

Quercus Margareta Ashe.

" *Marylandica* Muench.

" *minima* (Sarg.) Small.

" *minor* (Marsh) Sarg.

" *myrtifolia* Willd.

(Sand-hills.

Ware: Along Satilla River near Walbertown

Pierce: " " " NE of Waycross.

Charlton: Around Holkston, common

Charlton: Woods near Camden Co. line.

(Dry or rather dry pine-barrens.

Ware: Between Waycross and Rushin.

Glynn: Near Dock Junction.

Charlton: Dry pine-barrens around Holkston

Rather dry " " near Camp

Cornelia and Ellenton Mound (1577)

Thexia glabella flats of St. Mary's R.

Bugabo Island, Okefinokee Swamp.

Clinch: Near Fargo.

Charlton: Woods near Camden Co. line.

(Pierce: Driest sand-hills of
Satilla River (1466) (Photo)

Camden: Scrub thicket, Cumberland
Island (1540) (Photo)

Quercus nigra L.
(*Q. aquatica* Walt.)

Quercus Virginiana Mill.
(*Q. viciens* Ait.)

Quercus

Quercus

{ Along creeks and rivers.
Camden Near Woodbine
Charlton Folkston
Blinch Fargo

Pierce: Sand hills of Satilla River.

Glynn: Back of dunes, St. Simon's I.
Near Brunswick.

Camden: Cumberland; near Woodbine.

Charlton: West of Spanish Creek,
and near Traders Mill.

Blinch: Along Suwannee River, Fargo.

Echols: " Suwanoochee Creek.

Dry pine-barrens.

Ware: Vicinity of Braganza

Pierce: " " Mt. Mudge;
Between Schatterville and Hoboken.

Camden: Colesburg; South of Kingsland

Charlton: Folkston and vicinity.

Glynn: Dry pine-barrens
near Dock Junction (1576)

ULMACEAE

*Ulmus**Planera aquatica* (Walt.) Smel.*Celtis*

MORACEAE

Morus rubra L.

URTICACEAE

Boehmeria cylindrica (L.) Willd.

LORANTHACEAE

Phoradendron flavescens (Pursh) Kunt.Charlton: Along river at Traders Hill.
Swamp near Camden Co. line.Clinch: Muddy swamp of Suwannee
River near Fargo.Charlton: Woods near Camden Co. line.Charlton: Wooded bluff at Traders Hill.
Woods near Camden Co. line.

Moist places south of Valdosta.

Parasitic on trees, especially *Xyssa biflora*.
Charlton: Creek swamps east and
west of Goldston.Clinch: Suwannee River swamp, Fargo.

Eriogonum tomentosum Mx.

Polygonum arifolium L.

"

" *maritimum* L.

Polygonella

CHENOPODIACEAE

Chenopodium

Sand-hills and very dry pine-barrens.
Ware: Along Satilla R. near Walberton
and W. R. R.

Pierce: Sand-hills (ordinary) of
Satilla River. Vicinity of
Ft. Mudge, along B. & N. R. R.
near Schlatterville, from 44th
to 49th mile-post.

Charlton: Sand-hills near Folklon
and Traders Hill.

Echols: Along Allapaha River near Mayday
Always on the Columbia formation.
Not seen within 30 miles of the coast.

Camden: In Satilla River swamp
near Woodbine. (1563)

Camden: St. Mary's River swamp (1568)

Glynn: Drifting sands near
the beach, St. Simon's I. (1534)

Dry pine-barrens, Cumberland Island.

A weed. Brunswick, St. Simon's (1535).

Near Traders Hill; Bugaboo
Landing in Okefinokee Swamp.

Sondia

Salsola Kali L.

Salicornia ambigua Mx.

AMARANTHACEAE

Amaranthus

" *spinosus* L.

Alternanthera repens (L.) Runtze

Aenida

Froelichia Floridana (Vatt.) Moq.

BATIDACEAE

Batis maritima L.

{ Glynn: salt marsh, St. Simon's I.
Camden: inner shore, Cumberland I.

Glynn: Among wharves, Brunswick.
On dunes, St. Simon's Island.

Camden: " Cumberland "

Glynn: salt marshes, Brunswick
and St. Simon's Island.

Among the wharves, Brunswick.

" " " "

Streets of Brunswick.

Camden: Brackish marshes of
White Oak (1556), Satilla and
St. Mary's Rivers and Big Bayfish Br.

Charlton: Dry sandy cultivated fields
near Holkston and Traders Hill (1564)

Glynn: Salt marshes, Brunswick.

PHYTOLACCACEAE

Phytolacca decandra L.

NYCTAGINACEAE

Boerhaavia erecta L.

AIZOACEAE

Mollugo verticillata L.*Lescurium Portulacastrum* L.

"

PORTULACACEAE

Portulaca oleracea L." *pilosa* L.

CARYOPHYLLACEAE

Arenaria lanuginosa (Mx.) Rohrb.*Paronychia Baldwinii* (G. & H.) Chapm.*Stipulicida setacea* Mx.

Among ballast, Brunswick.

Wharves and streets of ...

Glynn: Brunswick } A weed.
Camden: White Oak }

Moist hollows of dunes, Cumberland I.

On ballast, Brunswick (1524)

{ Glynn: On ballast, Brunswick.
Camden: Around dwelling, White Oak.Ware: Railroad yard, Waveross.
Camden: Around dwelling, White Oak.

Under trees back of dunes, St. Simon's I.

" " " " " " " "

Pierce: Sand-hills of Satilla River.

NYMPHAEACEAE

Brasenia purpurea (Mx.) Caspary

Castalia

RANUNCULACEAE

Clematis

MAGNOLIACEAE

Magnolia glauca L.

" *grandiflora* L.

Liriodendron Tulipifera L.

Ware: Cypress pond between Waycross and Ruskin.

Ware: Between Ruskin and Glenmore.
Camden: Little Catfish Creek.
Charlton: Chase Prairie (Kefauver Swamp)
Schols: Between Haydon and Hayday
Lowndes: Cypress swamps S. of Valdosta.

Charlton: *Taxodium distichum* swamp about 4 m. N.E. of Holston.

Wet woods and non-alluvial swamps.
Ware, Pierce, Glynn, Camden,
Charlton, Schols, Lowndes.
East to within 15 miles of Brunswick.

Low rich woods.
Glynn: 7 or 8 miles from Brunswick.
Camden: Near Collesburg and Kingland.
Charlton: Traders Hill.
Clinch: Near Suwannee Creek.
Low woods along streams
Glynn: 7 or 8 miles from Brunswick
Camden: Little Catfish Creek.
Charlton: Near Holston & Traders Hill

ANONACEAE

Asimina angustifolia Gray." *parviflora* (Mx.) Bunal" *speciosa* Nash

LAURACEAE

Persea pubescens (Pursh) Sarg.*Sassafras officinale*

PAPAVERACEAE

Argemone alba

{ Dry pine-barrens.
Glynn: Near Dock Junction.
Camden: Cumberland Island
Charlton: Near Folkston

{ Dry or rich woods.
Camden: Woodbine
Charlton: Bluff at Traders Hill.

{ Dry pine-barrens.
Pierce: Between Hoboken and Schlatterville
Camden: Cumberland Island.
Charlton: Folkston and vicinity.

{ Wet pine barrens and non-alluvial swamp.
Ware: Near Glenmore (1470)
Camden: Sphagnum swamp of St. Mary's R.
Charlton: Okefinokee Swamp;
 Near Moniac and Ellicott's Mound.

Wooded bluff at Traders Hill.

Waste places, St. Simons I. (1880)

CRUCIFERAE

Warea cuneifolia (Muhl.) Nutt.

CAPPARIDACEAE

Polanisia tenuifolia T. & G.

SARRACENIACEAE

Sarracenia flava L.

Sarracenia minor Walt.

(*S. lacunosa* Barts., *S. variolans* Mx.)

Sarracenia psittacina Mx.

Pierce: Sand-hills of Satilla River,
perhaps not indigenous. (1465)

With the preceding (1467)

Moist pine-barrens.

Ware: Between Murrays and Beach.
Near Ruskin (rare)

Not seen within 60 miles of the coast.

Rather dry, moist or wet pine-
barrens or non-alluvial swamps.

Ware: Between Elsie and Upchurch; Near Waycross.

Pierce: Near Draganzza.

Camden: Sphagnum swamp of St. Mary's R.;
also common in pine-barrens.

Charlton: Common nearly everywhere,
especially in Okefinokee Swamp,
where it is 2-4 ft. tall. (1476)

Clinch: Near Fargo.

Schols: " "

Lowndes: Valdosta, and northward

Charlton: Moist pine-barrens between
Spanish Creek and Camp Cornelia.
Sphagnum bog, Okefinokee Swamp (1478)

DROSERACEAE

*Drosera**Drosera intermedia* Hayne.

SAXIFRAGACEAE

Decumaria Barbara L.*Itea Virginica* L.

HAMAMELIDACEAE

Hamamelis Virginiana L.*Liquidambar styraciflua* L.

Moist pine-barrens and sandy bogs.

Ware: Around shallow pond between

Ware and Ruskin.

Charlton: Vicinity of Holkiston and
Ellicott's Mound.

Bog near Bugaboo Landing, Chefinokee (1877)

Camden: Wet woods along and
near Little Catfish Creek.

Camden: Swamp of Little Catfish Cr. (1866)

Charlton: Swamp near Ellicott's Mound;
Chefinokee Swamp.

Echols: Near Suwannee Creek.

Pierce: Driest sand-hills of Satilla River.

Charlton: Sand-hills west of Spanish Cr.

Woods and swamps.

Ware: Along Satilla R. near Wattertown

Glynn: Near Dock Junction, & west.

Camden: Frequent.

Charlton: Along river at Traders Bluff;

Woods near Camden Co. Ga.

Blinch: Along Suwannee River.

ROSACEAE

Rubus cuneifolius Pursh.

" *nigrobaccus* Bailey.

Aronia arbutifolia (L.) Ell.

Brataegus apiifolia (Marsh.) Mx.

"

"

Chrysobalanus oblongifolius Mx.

Prunus

roadsides, etc.

Camden: West of Woodbine.

Charlton: Northeast Folkston.

Lowness: Valdosta.

Camden: Wet woods near cool spring south of King.

Moist pine-barrens and swamps.

Way: Near Glenmore.

Camden: West of Woodbine.

Charlton: Near Ellicott's Mound.

Okefinokee Swamp.

Along river at Traders Hill.

Sand-hills near " "

Charlton: Woods near Camden Co. line.

Sand-hills and very dry pine-barrens.

Pierce: Near Satilla River.

Charlton: Near Folkston, and on ridge near Camp Cornelia.

Wooded bluff at Traders-Hill.

LEGUMINOSAE

Mimosa strigillosa T. & G.*Morongia**Cercis Canadensis* L.*Bassia Marilandica* L.*Bassia occidentalis* L." *Tora* L.*Chamaecrista**Baptisia lanceolata* (Walt.) Ell.Charlton: Along St. Mary's River
Traders Hill (1503); Pine-barrens
northeast of Folkston.

Dry pine-barrens.

Camden: Cumberland Island; South
of Kingsland.

Along river at Traders Hill.

Charlton: *Toxodum distichum* swamp
about 6 m. northeast of Folkston

A weed. Brunswick.

{ " " , St. Simon's,
Folkston, ValdostaCharlton: Sand-hills southwest of
Folkston.

Dry pine-barrens.

Ware: North and west of WaycrossCamden: Near Vavely.Charlton: " Rappond & Folkston.

Baptisia perfoliata (L.) R. Br.

Erotaria Purshii DC.

" *rotundifolia* (Walt.) Poir.

Ruhriera pinnata (Walt.) Kuntze

Psoralea canescens Mx.

Amorpha *canadensis*

" *herbacea* Walt.

Petalostemon albidus (T. & G.)

" *carneus* Mx.

Ware: Along railroad between Waycross and Satilla River, perhaps not native.

Ware: Rather dry pine-barrens between Waycross and ~~Rossville~~.

Glynn: Rather dry pine-barrens between Brunswick and Dock Junction.

Sand-hills.

Charlton: Near Folkston.

Barlow: Along Allapaha River near Mayday.

Dry pine-barrens and weak sand-hills.

Camden: Cumberland Island.

Charlton: Near Folkston.

River-banks.

Camden: About a mile west of Woodbine.

Charlton: Traders Hill.

Charlton: Dry pine barrens - near Folkston.

Camden: Dry pine-barrens, Colesburg.

{ With the preceding, also
Charlton: Frequent around Folkston

Indigofera caroliniana Walt.

Bracca chrysophylla (Pursh) Kuntze

" *Virginiana* L.

Elodea vesicularis (Jacq.) Mohr

Aeschynomene viscidula Mx.

Stylosanthes biflora (L.) B.S.P.

Gomphia bracteata (Walt.) Griseb.

{ Dry pine-barrens and sand-hills.
Camden: Cumberland Island.
Charlton: West of Spanish Creek

Charlton: Sand-hills between Spanish Creek and Traders Hill (1497)

{ Dry pine-barrens and sand-hills.
Ware: Along Satilla River near Waltertown.
Pierce: " " " " Also near Ft. Mudge.
Charlton: Near Racepoint, Folkston and Butler.

Not seen in the maritime counties.

{ Apparently a weed.
Camden: Rice plantation W. of Woodbine.
Charlton: South & west of Folkston.

{ Glynn: Streets of Brunswick.
Camden: Dry pine-barrens, Cumberland I.
Charlton: Sand-hills near Traders Hill (1498).

Camden: Dry pine-barrens, Cumberland I.
Charlton: Sand-hills around Folkston.

Dry pine-barrens, Cumberland Island.

Meibomia arenicola Vail.

Lespedeza striata (Thunb.) ...

Clitoria Mariana L.

Apios tuberosa Moench.

Galactia Elliottii Nutt.

Galactia volubilis (L.) Britton

Dolicholus simplicifolius (Walt.) Vail.

Charlton: Sand-hills S.W. of Folkston

A weed. Near Brunswick and Vidalia.

Dry pine-barrens, Cumberland Island.

Camden: Swamp west of Bull Head Bluff.

Rather dry pine-barrens.

Glynn: East and west of Dock Jc.

Camden: Cumberland Island; White Oak.

N. of Colerburg (1559); S. of Kingsland.

Charlton: Folkston to Camp Cornelia.

Bugabo Island (perhaps introduced)

Blinch: Near Fargo (rare)

Not seen more than 70 miles inland.

Woods near White Oak.

Dry pine-barrens and sand-hills.

Glynn: Near Dock Junction.

Camden: South of Kingsland

Charlton: Near Spanish Creek.

LINACEAE

Linum floridanum (Pland.) Thel.

ZYGOPHYLLACEAE

Gribulus cistoides L.*Hallstroemia maxima* (L.) F. & A.

RUTACEAE

Xanthoxylum blava-herculis L.

POLYGALACEAE

Polygala cruciata L." *cymosa* Walt.

Rather dry pine - barrens, etc.

Ware: Between Bolen and Elsie. ^{Near} Waycross.

Charlton: Near Folkston, and westward

On ballast, Brunswick. (1522)

Streets of Brunswick (1524)

Among dunes, St. Simon's and Cumberland

Moist pine-barrens.

Ware: Between Waycross and river.

Camden: Colesburg

Charlton: East side of ridge; Moniac

Pine-barren ponds.

Ware: North of Durrys; Betw. Beach & Bolen.

Between Waycross & Rushin; Near Elmore.

" " " Schlatterville.

Wayne: West of Kahunta

Camden: Woodbine; South of Ringland.

Charlton: North & south of Camp Conelia.

Clich: East of Fargo; Near Shelma

Polygala grandiflora Walt.

" *incarnata* L.

" *lutea* L.

" *ramosa* Ell.

" *setacea* Mx.

Polygala

Charlton: sand-hills near Folkston.

Ware: Moist pine-
Waycross and Satilla River.

{ Moist pine-barrens.

Ware: Northeast & southwest
Glynn: Northwest of Cock Is. Waycross

Camden: Wood?

Charlton: Near Folkston; East slope
of Pinckney Ridge.

{ Moist pine-barrens, rather common.
Ware, Camden, Charlton.

{ Rather dry pine-barrens.

Ware: Between Waycross & Ruskin.

Camden: Wood?

Charlton: Spanish Creek to Camp
Near Ellicott's Mound.

Blinch: " Fargo.

{ Rather dry pine-barrens.

Ware: Near Waycross (1468)

Camden: West of Bluff.

Charlton: Spanish Creek to Okefinokee
Northeast of Folkston.

EUPHORBIACEAE

Croton argyranthemus Mx." *glandulosus septentrionalis* Willd. Ag." *maritimus* Walt.*Crotonopsis spinosa* Nash.*Jatropha stimulosa* Mx.*Stillingia sylvatica* L.*Sebastiana ligustrina* (Mx.) Willd. Ag.Charlton: Sand-hills around Folkeston.

A weed. Brunswick and Vicksburg.

{ Glynn: Dunes, St. Simon's Island.
Camden: Beach, dunes and inner shore, Cumberland Island.Charlton: *Rhexia glabella* flats near St. Mary's.{ Charlton: Sand-hills near Folkeston.
Camden: Dry pine-barrens, Cumberland.{ Dry pine-barrens or sand-hills.
Ware: Between Waycross and Dublin.
Camden: Cumberland Island;
South of Kingland.
Charlton: Near Folkeston.Charlton: Along St. Mary's River at Frederic Hills.

Euphorbia cordifolia Ell.

" *gracilis* Ell.

" *maculata* L.

" *polygonifolia* L.

" *...*

"

ANACARDIACEAE

Rhus copallina L.

" *radicans* L.

Dry pine-barrens, Cumberland Island.

Charlton: Sand-hills west of Spanish R.

{ A weed, growing especially on railroads.
Brunswick, White Oak, Kolesburg,
Camp Korrin, Moniac, Valdosta.

{ Beach, St. Simon's Island.
Dunes, Cumberland "

Glynn: Along railroad near Dock Junction,
(5 feet tall)

On ballast, Brunswick. (1520)

Pierce: Driest sand-hills of Satilla R.

Glynn: Dry pine-barrens near Dock Jc.

Camden: " " " south of Woodbine.

Charlton: West of Spanish Creek;
Wooded bluff at Traders Hill.

Camden: St. Marys river swamp; Wet woods.

Charlton: Swamp east of Charlton.

Chefinokee Swamp, not common.

Rhus Toxicodendron L.

" *Vernix* L.

CYRILLACEAE

Cliftonia monophylla (Lam.) Button

Cyrilla racemiflora L.

ILICACEAE

Ilex Caroliniana (Walt.) Treb.

" *Bassine* L.

Pierce: Sand-hills of Satilla River.

Wet woods and non-alluvial swamps.

Charlton: Near Golston and Traders Hill.

Lowndes: South of Valdosta.

{ Swamps. Ware: Near Wattertown.
Camden: West of Bull Head Bluff.

{ Swamps, rarely associated with the preceding.

Ware: Along Satilla R. near Wattertown.

Pierce: West of Schlatterville.

Charlton: Near Golston, Traders Hill
and Moniac; Okefinokee Swamp.

Clinch: Near Fargo and Shelma.

Echols: " Suwanoochee Creek.

Lowndes: Near Blanton.

Pierce: Sand-hills of Satilla River.

{ Camden: Wet woods south of Ringland.

Charlton: River swamp near Moniac.
Okefinokee Swamp.

Olex glabra (L.) Gray

" *lucida* (Ait.) T. & G.

" *myrtifolia* Walt.

" *opaca* Ait.

" *vomitaria* Ait.

ACERACEAE

Acer rubrum

From rather dry pine-barrens to swamps.

Glynn: Just back of Brunswick.

Camden: Woodbine; St. Mary's R. swamp.

Charlton: Near Folkston & Ellicott's mound.

Okefinokee Swamp, near Fargo.

Clinch: Near Fargo.

Wet pine-barrens, sandy swamps, etc.

Ware: Near Glenmore.

Charlton: " Folkston; Okefinokee Swamp.

Ware: Cypress pond between Waycross & Rushin.

Charlton: Similar localities near Butler,

Moniac and Ellicott's mound.

Clinch: Suwannee River swamp.

Cypress swamp near Headlight.

Echols: Near Mayday.

Not seen within 40 miles of the coast.

Camden: Swamp near Bull Head Bluff.

Charlton: St. Mary's River swamp, Moniac.

Camden: " " " "

Wet woods and swamps, frequent.

Glynn, Camden, Charlton, Clinch

Echols, Lowndes. Not near salt water.

RHAMNACEAE

Berchemia scandens (Hill.) Trel.*Ceanothus microphyllus* Mx.

VITACEAE

Vitis aestivalis Mx." *rotundifolia* Mx.*Ampelopsis arborea* (L.) Rusby.

Camden: Swamp of Little Catfish Creek.
Charlton: Along river at Fredericks Hill.
 Swamp about 6 m. N.E. of Fredericks.

Camden: Dry pine-lands south
 of Kingsland.

Camden: Near Satilla River at Kingsland.
Charlton: Wooded bluff at Fredericks Hill.

Glynn: Between Dock Jc. and Barton.
Camden: Swamp west of Bull Head
 Bluff; On dunes, Cumberland I.;
 Wet woods south of Kingsland.
Charlton: Sand-hills west of Spanish Cr.
 Wooded bluff at Fredericks Hill.

Along canal, Okefinokee Swamp,
 introduced.

Glynn: Dunes, St. Simon's Island.
 Woods near Dock Junction.
Charlton: Woods near Camden Co. line.

Panthenocissus quinquefolia (L.) Planch

MALVACEAE

Sida rhombifolia L.

Kosteletzkya Virginica (L.) A. N. S. P.

Hibiscus aculeatus Walt.

THEACEAE

Gordonia Lasianthus L.

STERCULIACEAE

Melochia corchorifolia L.

In woods.

Camden: Near White Oak, Satilla
River and Big Catfish Creek.

Charlton: Near Traders Mill.

A weed. Brunswick and Valdosta.

Camden: Brackish marshes of White
Oak, Satilla (1565) and St. Mary's.

Rather dry pine-barrens.

Charlton: Between Spanish Creek
and Chefinokee Ridge.

Lowndes: South of Valdosta.

Camden: St. Mary's River swamp (1571)

Charlton: Non-alluvial swamps near
Folkston, Moniac and Ellicott's Road
Chefinokee Swamp.

In cultivated ground near
dwelling, White Oak

HYPERICACEAE

Ascyrum humilum Michx." *stans* Michx.*Hypericum*" *fasciculatum* Lam." *galioides* Lam." *mutilum* L.

{ Rather dry pine - barrens.
 Ware: Between Ware and Ashkin.
 Charlton: Near Folkston.

{ Rather dry or moist - barrens.
 Hlynn: Between Spanish Brook and Rock Jc.
 Camden: Bolesburg.
 Clinch: Near Fargo.

Dry pine-barrens, Cumberland I.

Moist sandy places along railroad
just south of Folkston.Moist pine-barrens, frequent.
Ware, Camden, Charlton, Clinch, Lowndes.Charlton: Along river at Traders Hill
(1502); " Spanish Brook.Low grounds near plantation west
of Woodbine, Camden Co.

Hypericum myrtifolium Lam.

"

"

" *spacum* F. & G.

" *pilosum* Walt.

"

Sarothra gentianoides L.

Triadenum Virginicum (L.) Raf.

Moist pine-barrens, frequent.
Ware, Blynn, Camden, Charlton.

Charlton: Along river at Traders Hill.
(1501); " Spanish Creek.

Clinch: Along Suwannee River near Fargo.

Charlton: Mixed with no. 1501.)

Moist pine-barrens,
Ware, Camden, Charlton, Clinch.

Camden: Moist pine-barrens, Colesburg.

Ware: Wet pine-barrens between
Ruskin and Glenwood.

{ Rather dry pine-barrens; mostly ruderal.
Pierce: West of Schlatterville.
Camden: Between Woodbine & Colesburg.
Charlton: East of Folkston.
Echols: Mayday, and westward.

Charlton: Okelake Swamp.

Clinch: Wet pine-barrens east
of Fargo.

CISTACEAE

*Helianthemum**Lechea*

"

VIOLACEAE

Viola denticulosa Pollard.

TURNERACEAE

Piriqueta Caroliniana

PASSIFLORACEAE

Passiflora lutea L.

CACTACEAE

Opuntia vulgaris Mill.

Dry sand east of P. R. Cumberland Co.

{ Rather dry pine-barrens, of - around ponds.
 Ware: Between Waycross and Rush.
 Charlton: North and south of Camp Louisa.
 Not seen east of Okefenokee Ridge.

Ware: Dry sand between M.
 and Satilla River.

Charlton: *Taxodium distichum* swamp
 about 4 miles northeast of P. R.

Charlton: Along O. & P. R. R. northeast
 of Folkston.

{ Apparently ruderal.
 Camden: Near house west of
 Bull Bluff; Cumberland Co.

Sand-hills, etc.

Ware: Between Waycross and -

Pierce: Along Satilla River.

Charlton: Near Tradewind.

Opuntia *Pes-borvi*, Leconte.

LYTHRACEAE

Decodon verticillatus (L.) Ell.

Lythrum lineare L.

"

MELASTOMACEAE

Rhexia ciliosa Mx.

" *glabella* Mx.

"

" *lutea* Walt.

{ Glynn: Back of house, Ht. ...
Camden: Dry sandy, Cumberland "

Along canal, Okefinokee Swamp.

Camden: Brackish marshes of
 W. River (1553)

Charlton: *Taxodium distichum*
 swamp - near Camden Co. line.

{ Moist pine-barrens, nowhere ant.
 { Ware, Camden, Charlton, Clinch, Lowndes.

{ Moist pine-barrens, etc., usually
 very abundant. The predominating
 species in the "Rhexia glabella"
 { Ware, Pierce, Glynn, Camden,
 Charlton, Clinch, Lowndes.

Moist pine-barrens, ...

Ware: Near Rushin

Camden: Around small ponds south
 of Big Batfish Creek.

Charlton: *Rhexia glabella* - flats of
 ... Creek.

Rhexia

" *floridana* Nutt.

" *serrulata* Nutt.

" ?

ONAGRACEAE

Oenardia palustris L.

Ludwigia alata Ell.

" *capitata* Mx.

{ Moist pine barrens and swamp ponds.
Ware: East of Kingsland.
Camden: South Kingsland.

Camden: Margin of shallow pond south of Kingsland. (1572)
Clinch: Muddy flats in Lawrence River swamp (1578)

Camden: Moist pine-barrens near Woodbine.

{ Wet pine-barrens, rather common.
Ware, Pierce, Glynn, Charlton, Lowndes.

Camden: Edge of Satilla River marsh, Woodbine; Swamp of Little Catfish Creek.

Camden: Brackish marshes of White Oak (1554), Satilla & St. Mary's Rivers.

Moist pine-barrens.
Charlton: East-west (1488) of ridge near Camp Cornelia; East of Folkston.
Okefinokee Swamp (peculiar form, 1485)
Clinch: East of Kingsland.

Ludwigia alternifolia L.

" *hirtella* Raf.

" *linearis* Walt.

" *linifolia* Poir.

" *pilosa* Walt.

" *sphaerocarpa* Ell.

" *virgata*
linearis (1887)

Lowndes: ditches along railroad south
 of Vald.

Rather dry pine-barrens,
Camden; Colesburg;
Lowndes: Valdosta.

{ Ditches, moist pine-barrens, etc.
Camden: Colesburg; south of
Lowndes: South of Valdosta

Charlton: *Taxodium imbricarium*
 swamp near St. Mary's River
 southeast of

Wet pine-barrens, shallow ponds, etc.
Ware, Glynn, Camden, Charlton,
Clinch, Echols, Lowndes. Extends
 to within 12 miles of Brunswick.

Ditches, Valdosta.

Charlton: Ground cypress and
 near St. Mary's River east
 of Ellicott's Ground.

Ludwigia

Jussiaea decurrens (Walt.) Bb.

" *leptocarpa* Nutt.

Oenothera humifusa Nutt.

Gaura angustifolia Mx.

HALORAGIDACEAE.

Proserpinaca palustris L.

" *pectinata* Lam.

Ayriophyllum laxum Nutt.

Camden: between dunes and beach, Cumberland I. (1542)
Rather dry pine - west of Woodbine and south of Kingsland.
Charlton: ?
Lowndes: S. of V

{ Ditches and low grounds.
Camden: Plantation west of Woodbine.
Lowndes: Near Valdosta.

" " "

Meadow near beach, Cumberland I.

{ Blynn: Brunswick } Apparently ruderal
Camden: Cumberland
Charlton: Swamp near Camden Co. Ga.

Camden: West of Bull Head Bluff.

Camden: Shallow pond south of Kingsland.

In canal, Okefenokee Swamp.

ARALIACEAE
Aralia spinosa L.

UMBELLIFERAE
Hydrocotyle umbellata L.

Pentstemon Asiatica (L.) Urban

Eryngium aromaticum Baldw.

" *Baldwinii* Spreng.
 (E. gracile Baldw.)

Rich woods.

Glynn: Near Dock Jc. and Carters.

Camden: Near Satilla R. at W. 1st St.

Between Big & Little Satfish Creeks.

Charlton: Bluff. Traders Mill.

Blinch: Near Suwannee Creek.

Glynn: Artesian well, St. Simon's.

Camden: Inner shore, Cumberland.

Charlton: Okefenokee Swamp (1471)

Lowndes: Ditches south of Valdosta.

Wet pine-barrens and ponds.

Ware: Between Waycross & Rushin.

Glynn: " Brunswick & Dock St.

Camden: Woodbine; south of Kingsland.

Charlton: Frequent.

Charlton: Sand-hills (1496) and dry
 pine-barrens from Folkston to
 Camp Cornelia.

Blinch: Near Fargo (not common)

Charlton: Dry pine-barrens near
 Folkston (1494); along river at
 Traders Mill.

Eryngium Ludovicianum Morong.

" *synchaetum* (Gray) B. & R.

" *virgatum* Lam.

Cicuta Burtissii B. & R.

Oxypholis filiformis (Walt.) Britton

Charlton: Moist pine-barrens about
5 miles northeast of Folkston.

{ Rather dry pine-barrens, etc.

Ware: Near Waycross.

Camden: West of Bull Head Bluff;

Bolesburg; Near St. Mary's River.

Charlton: *Rhexia glabella* Blats.

Near Folkston, Camp Cornelia,
and Ellicott's Mound.

Blinch: Near Fargo.

{ Moist pine-barrens.

Charlton: Moniac.

Lowndes: Valdosta (abundant)

{ Camden: Swamp about a mile
west of Bull Head Bluff.

Brackish marshes of White Oak River.

Moist pine-barrens and shallow ponds.
Seen in every county except Wayne
(but doubtless occurs there too).

Extends east to within 10 miles
of Brunswick.

CORNACEAE

Nyssa biflora Walt." *Ogeche* Marsh." *uniflora* Wang.

CLETHRACEAE

Clethra

ERICACEAE

Bejaria racemosa Vent.

Pine-baren ponds and swamps.
 Ware, Glynn, Camden, Charlton,
 Echols, Lowndes. Rare in Shefina
 Swamp. Extends east to within
 12 miles of Brunswick.

In rivers, creeks, branches and swamps.
 Ware, Pierce, Camden (Bull Head Bluff),
 Charlton, Clinch, Echols. Not
 seen within 20 miles of the coast.

Glynn: Swamps between Dock Junction
 and Cartier, Bladen and S.B.F.
 Crossing, and southward.

Swamps and wet pine-barens.
 Ware, Pierce, Camden (Bull Head Bluff),
 Charlton, Clinch, Lowndes. Not
 seen within 20 miles of the coast.

Dry or rather dry pine-barens.
 Camden: Cumberland Island (1539)
 Woodbine; Colesburg; South of Kingsland.
 Charlton: Frequent between Butler
 and Monice; Northwest of
 Ellicott's Mound.

Azalea viscosa L.

Kalmia hirsuta Walt.

Leucothoe

Pieris nitida (Bart.) B. & Z.

Pieris phillyreaefolia (Hook.) DC.

Xolisma

{ Swamps, etc.

Ware: Vicinity of Braganza

Charlton: Just west of Folkston.

Echols: Near Suwannee River.

{ Rather dry pine-barrens.

Camden: Woodbine.

Charlton: Near Camp Cornelia;
Bugaboo Island.

{ Charlton: In Okefinokee Swamp.

Blinch: West bank of Suwannee River.

{ Non-alluvial swamps, etc.

Ware: Near Glenmore.

Camden: Cumberland (1541); Woodbine;
Sphagnum swamp of St. Mary's River.

Charlton: Vicinity of Folkston & Moniac.
Okefinokee Swamp.

Charlton: Cypress pond north of
Camp Cornelia; Okefinokee Swamp (1475, 1477)

Camden: Moist pine-barrens
south of Kingsland.

Xolisma

VACCINIACEAE

Gaylussacia

" *dumosa* (Andr.) F. & G.

Vaccinium nitidum Andr.

Rather dry pine-barrens and sand-hills.
Ware: Between Waycross and Ruskin.
Pierce: Dry sand-hills of Satilla R.
 (abundant); Vicinity of Ft. Mudge.
Glynn: Vicinity of Dock Junction.
Camden: Woodbine, south of Kingsland.
Charlton: Bugaboo Island; and
 frequent outside of the swamp.
Clinch: Near Fargo.

Rather dry pine-barrens.
Glynn: Near Dock Junction.
Camden: South of Kingsland.
Charlton: Bugaboo Island.

Dry or rather dry pine-barrens.
Charlton: Near Folkston; Bugaboo I.

Dry or rather dry pine-barrens.
Glynn: Vicinity of Dock Junction.
Camden: South of Little Catfish Cr.
Charlton: Near Folkston, Butler, &
 Ellicott's Mound; Bugaboo Island.
Clinch: Near Fargo.

Batodendron arboreum Nutt.

Polycodium stamineum (L.) Greene

PRIMULACEAE

Samolus floribundus L.B.K.

Anagallis arvensis L.

SAPOTACEAE

Bumelia

" *lycioides* (L.) Pers.

Sand-hills and rich woods.

Pierce: Sand-hills of Satilla River.

Camden: Near Satilla River, Woodbine.

Charlton: Sand-hills west of Spanish Br.

Wooded bluff at Traders Hill.

Clincks: Woods near Suwannee Creek.

Sand-hills and very dry pine-barrens.

Pierce: Driest sand-hills of Satilla River (6 ft. tall)

Charlton: Near Raceford & Holston.

Camden: Brackish marshes of White Oak River; Wetwoods south of Kingsland.

On ballast, Brunswick.

Glynn: Back of dunes, St. Simon's Island

Camden: Dry pine-barrens, Cumberland I.

Charlton: Wooded bluff at Traders Hill

Charlton: Along river at Traders Hill.

EBENACEAE

Diospyros Virginiana L.

OLEACEAE

Fraxinus Caroliniana Mill.*Osmanthus Americana* (L.) A. N. S.

LOGANIACEAE

Gelsemium sempervirens (L.) Ait. f.*Polypremum procumbens* L.

Habitat very variable.

Pierce: Sand-hills of Satilla River.Hlynn: About 10 miles from ...Camden: Woods, White Oak.Charlton: Frequent, from sand-hills to swamps.

Swamps.

Camden: West of Bull Head Bluff.Charlton: Along Spanish Creek and St. Mary's River.Blinch: Suwannee " swamp.Echols: Suwannee Creek " "

Sand-hills and rich woods.

Pierce: Driest sand-hills (fruiting)Charlton: West of Spanish Creek.Blinch: Woods near Suwannee Cr.

In woods.

Camden: Near White Oak; S. of KingslandCharlton: Bluff at Traders Hill

A weed, or mostly so.

Charlton: Folkston.Camden: Inner shore, Cumberland I.Echols: Near Howell.Lowndes: " Blanton

Gynostomum sessilifolium (Walt.) Griseb.

GENTIANACEAE

Sabbatia campanulata (L.) Torr.

" *decandra* (Walt.) Harper.

" *Elliottii* Steud.

" *foliosa* Fernald

" *gentianoides* Ell.

" *macrophylla* Hook.

" *stellaris* Pursh.

Charlton: Moist pine-barrens near
Chefinslee Ridge and Elliott's Road.

Charlton: *Rhexia glabella* flats of
St. Mary's River and Spanish Cr.

{ Pine-barren ponds, etc.
Ware: Near Waycross, Ruskin and
Glenmore.
Charlton: East slope of Chefinslee Ridge.

Rather dry pine-barrens, common.
Glynn (to within 10 miles of Brunswick),
Camden, Charlton, Blinch, Echols, Lowndes.

{ Ware: Along Satilla R. near Waycross.
Camden: Swamp of Little Catfish Cr.
Charlton: Along river at Traders Hill.

Moist pine-barrens.
Charlton: Several stations east
and west of Folkston.

Charlton: Moist pine-barrens near Uptonville.

{ Glynn: Around artesian well, St. Simon's I.
Camden: Meadow near beach, Cumberland.

Portonia

Limnanthemum aquaticum (Walt.) Britton

APOCYNACEAE

Amsonia ciliata filifolia Wood.

ASCLEPIADACEAE

Asclepias cinerea Walt.

" *humistrata* Walt.

" *lanceolata* Walt.

" *perennis* Walt.

Charlton: Sphagnum bog about
2 miles east of Folkston.

Okef: Swamp.

Pierce: Sand-hills of Satilla River.

{ Rather dry pine-barrens.
Camden: Colesburg.
Charlton: Frequent between Folkston
and Okefinokee Ridge.

Charlton: Sand-hills between
Spanish Creek and Traders Bluffs.

{ Wet pine-barrens and ponds.
Ware: Between Waycross and Ruskin
Pierce: " Draganya & Ft. Mudge.
Glynn: Near Thalman; south of
S. B. T. Crossing.
Camden: Near Waverly
Charlton: Around Folkston.

Charlton: Bank of St. Mary's River
at Traders Bluffs

Anantherix connivens (Baldw.) Gray

Podostigma pubescens Ell.

CONVOLVULACEAE

Sporobolus filicanalis Blume.

" *sagittata* Poir.

Breweria humistrata (Walt.) Gray

CUSCUTACEAE

Cuscuta compacta Juss.

BORAGINACEAE

Heliotropium curassavicum L.

" *Europaeum* L

{ Moist pine-barrens.

Charlton: North of Camp Cornelia.

Lowndes: Valdosta.

{ Not seen east of ridge (but discovered in Camden Co. by Dr. Baldwin)

Rather dry pine-barrens.

Camden: Woodbine (1 specimen); also

St. Mary's (according to Elliott)

On ballast, Brunswick (1523)

Charlton: 5 miles northeast of Folkston.

Camden: Dry pine-barrens, Cumberland &

Charlton: Sand-hills S.W. of F. 1524.

{ On shrubs in swamps.

Camden: South of Woodbine.

Charlton: Near Folkston; Okefinokee

Swamp, along canal.

On ballast, Brunswick. (1525)

" " " (1526)

HYDROPHYLLACEAE

Nama corymbosa (Macbride) Kuntze

VERBENACEAE

Lippia nodiflora (L.) Mx.*Callicarpa Americana* L.

LABIATAE

Merosphaerum rugosum (L.) Pollard*Teucrium Nashii* Kearney*Physostegia*Charlton: Moist pine-barrens about 5 miles northeast of Folkston (1516)

{ Not far from the coast. Often a weed.

Glynn: Streets of Brunswick; Around artesian well, St. Simon's Island.Camden: Meadow near beach, Brunswick.Charlton: Moist pine-barrens about 5 miles northeast of Folkston (1513)

{ Habitat variable. Usually in dry places.

Glynn: Back of dunes, St. Simon's Island, near Thalman's.Camden: Near White Oak and Woodbine.Charlton: Sand-hills and wooded bluff at Traders Bluff.Clinch: Woods near Suwannee.{ Moist pine-barrens, margins of swamps, etc.
Glynn, Camden, Charlton, Lowndes.Charlton: First *Taxodium distichum* swamp east of Folkston (1510)Charlton: With the preceding; also near Ellicott's mound.

Monarda punctata L.

Lycopus

Roellia hyssopifolia (Benth.) Britton

" *nuda* (Nutt.) Kuntze

SOLANACEAE

Solanum carolinense L.

"

"

Satura latula L.

Nicotiana ?

Dry sandy woods, etc.

Blynn: Just back of Brunswick.

Camden: Woodhouse Station - Big and
Little - both - green.

Charlton: Swamp near Camden Co. Co.

{ Moist pine-barrens, etc.

Camden: W. of Bull Head Bluff; Colesburg.

Charlton: *Rhexia glabella* f. etc.

St. Mary's River and Spanish Creek.

Near Moniac.

Clinch: Southwest of F. g.

{ Moist pine-barrens.

Ware: Near Solen and Waycross.

Camden: Colesburg.

Charlton: West of Spanish Cr.; Near Moniac.

A weed. Brunswick.

On ballast, " (1518)

" " " (1519)

Charlton: Roadside west of Kolbston

Streets of Brunswick.

SCROPHULARIACEAE

Verbascum Thapsus L.*Monnina Monnina* (L.) Britton" *acuminata* (Walt.) Kunze*Scoparia dulcis* L.*Micranthemum orbiculatum* Mx.*Dratiola pilosa* Mx.*Sophronantha hispida* Benth.

A weed, Cumberland Co.

{ *Glynn*: Artesian well, St. Simon's I.
Camden: Inner shore,{ *Bl.*: Pine-barrens about 5 mi.
northeast of Folkston.{ A weed.
Glynn: Between Brunswick & Oak Jc.
Camden: West of Bull Head Bluff
Lowndes: ...{ *Camden*: Wet woods south of Kingsland.
Charlton: Along river & Traders Hill.{ Moist pine-barrens.
Charlton: N. W. of Ellicott's Mound.
Clinch: Southwest of Fargo.
Lowndes: Valdosta.{ Rather dry pine-barrens, etc.
Ware: Between Waycross and Ruskin
Pine: Sand-hills of Satilla River.
Charlton: Near Camp Cornelia & Traders Hill
Clinch: Southwest of Fargo.

Euchnera

Aegelia cassioides (Walt.) Gmel.

Sasytoma pectinata (Walt.) Benth.

" *Virginica* (L.) Britton

Gerardia fasciculata Ell.

" *linifolia* Walt.

BIGNONIACEAE

Bignonia crucigera L.

LENTIBULARIACEAE

Utricularia juncea Vahl.

" *purpurea* Walt.

Charlton: Rather dry pine-barrens
near Folklton.

{ Rather dry or moist pine-barrens.
Ware: Between Waycross and Ruskin.
Charlton: Along river near Camp Cornelia.
Northwest of Ellicott's Mound.

Charlton: Sand-hills southwest of Folklton

" " " west " "

Camden: Meadow between dunes
and beach, Cumberland I. (1543)

Charlton: Cypress pond south of
Camp Cornelia; Chefinokee Swamp.

{ Low woods and swamps.
Glynn: Near Dock Junction.
Charlton: Along Spanish Creek.
Echols: Near Suwannee " "

Charlton: Sphagnum bog east of Folklton.
Chefinokee Swamp, Glasse Prairie.

" " " "

ACANTHACEAE

Tubiflora Carolinensis (Walt.) Griseb.

PLANTAGINACEAE

Plantago sparsiflora Mx.

RUBIACEAE

Houstonia rotundifolia Mx.*Oldenlandia uniflora* L.*Diodia teres* Walt." *Virginiana* L.Camden: Moist pine barrens
west of Bull Head Bluff.Charlton: Rather dry pine barrens
about 5 m. northeast of Folkston (1512)

{ Dry pine - barrens, etc.

Glynn: Near Dock Junction.Camden: Cumberland Island.Charlton: Sand-hills east of Folkston.Blinch: Northwest of Fargo.

{ Rather dry pine-barrens.

Camden: Between Woodbine & Colerburg.Charlton: North of Camp Columbia.

Mostly a weed, especially along railroads.

Pierce, Glynn, Camden, Charlton,
Echols, Lowndes.

{ Usually in moist places, often a weed.

Camden: White Oak; Colerburg.Charlton: Moist pine-barrens near Moniac.

Richard

Pinckneya pubens Mx.

Cephalanthus occidentalis L.

Mitchella repens L.

Galium hirsutum Mx.

CAPRIFOLIACEAE

Sambucus canadensis L.

Viburnum molle Mx.

A weed. Valdosta.

{ Swamps and wet pine-barrens.
Pierce: East and west of Schlatterville.
Echols: Near Mayday.
Lowndes: " Blanton and Valdosta.
 Not seen east of ridge.

{ In swamps, frequent.
Ware, Camden, Charlton, (including
Cherokee Swamp), Clinch, Lowndes.

Camden: Wet woods south of
 Kingsland.

{ Camden: Dry pine-barrens; Cumberland
Charlton: Wooded bluff at Traders Hill

Camden: Satilla River swamp near Woodbine

Charlton: Bank of St. Mary's
 River at Traders Hill.

Viburnum nudum L.

" *obovatum* Walt.

Lonicera sempervirens L.

CUCURBITACEAE

Melothria pendula L.

LOBELIACEAE

Lobelia cardinalis L.

Wet pine-barrens and swamps.

Ware: Near Auckin.

Glynn: About 10 miles from Brunswick.
Between Bladen and S.B.T. crossing.

Camden: Colesburg.

Charlton: Between Folkston and
Bull Head Bluff.

Charlton: Near river at Traders Mill.

Pierce: Driest sand-hills of
Satilla River.

Glynn: Under trees just back of
dunes, St. Simon's Island.

Camden: Swamp of Little Catfish Cr.

CICHORIACEAE

Silene caroliniana (Walt.) Raf.

AMBROSIACEAE

Iva frutescens L." *imbricata* Walt." *microcephala* Nutt.*Ambrosia artemisiifolia* L." *psilostachya* DC.

Streets of Brunswick.

Glynn: Salt marshes around Brunswick.
Dunes, St. Simon's Island.

" " " "

A weed along B. & P. R. R. northeast
of Holkston, in Camden and Charlton.{ A weed. Less common in this region
than farther north.Glynn: Waste places, St. Simon's I.Camden: West of Bull Head Bluff.Charlton: Rather dry pine-barrens
about 5 miles north of Holkston
(1514). To all appearances indigenous.

8/

COMPOSITAE

Vernonia angustifolia Mx.

"

Elephantopus scatus Bertol.

" *Carolinianus* Willd.

"

Eupatorium album L.

Dry or rather dry pine-barrens.
Glynn, Camden, Charlton, Blinch.

{ Swamps, etc.
Camden: Near Woodbine.
Charlton: Southeast and northeast
of F. Charlton.

{ Dry or rather dry pine-barrens.
Glynn: Near Brunswick.
Camden: Cumberland Island.
Charlton: East of Folkston and
Camp Cornelia.

Charlton: Wooded bluff at Traders Bluff.

{ Rather dry pine-barrens.
Camden: North of Colesburg (1560);
South of Kingsland.
Blinch: Southwest of Fargo.

Pierce: Dry pine-barrens near Ft. Dodge.

Eupatorium capillifolium (Lam.) Small.

{ A weed, mostly in swamps.
Glynn: Just east of Cartier.
Charlton: Along B. & O. R. in swamp
 near Camden.
 Along canal in Okefenokee Swamp.

" *compositifolium* Walt.

{ In dry pine-lands, doubtfully natural.
Ware, Pierce, Camden (Bamberland
 Island), Echols (Howell).

" *hypsipifolium* L.

{ Dry pine-lands.
Glynn: Between Brunswick & Duck.
Camden: Woodbine.

"

Blinch: Muddy places in swamp
 of Suwannee River (1579)

" *perfoliatum* L.

{ Swamps.
Charlton: First *Saxodium distichum*
 swamp east of Folkston.
Glynn: Between Cartier and Thalmers.

" *rotundifolium* L.

{ Moist or rather dry pine-lands.
Ware, Pierce, Glynn, Camden,
Charlton, Blinch, Lowndes.

Eupatorium semiserratum Ell.

" *serotinum* Mx.

" *verbenaefolium* Mx.

Conoclinium coelestinum (L.) Ell.

Mikania scandens (L.) Willd.

Charlton: First *Exodium distichum*
swamp east of Folkston. (1511).

Camden: Satilla River swamp
opposite Woodbine.

{ Rather dry pine-barrens.
Camden: Woodbine.
Charlton: Near Folkston.
Lowndes: South of Valdosta.

Camden: Wet woods south of Kingsland
Charlton: Along river at Traders Hill.

Glynn: Ground artesian well, St. Simons.
Swamps between Dock & ... Barter.
" " Bladen & S. B. T. crossing

Camden: In nearly every swamp
along the railroad, from Waverly
to St. Mary's River (1567)

Charlton: Along river at Traders Hill.
Swamp east of Folkston.

Lilisa odoratissima (Walt.) Cass.

" *paniculata* (Walt.) Cass.

Carphophorus

Heterotheca subaxillaris (Lam.) Britton & S. P.

Chrysopsis graminifolia (Mx.) Nutt.

"

{ Rather dry or moist pine-barrens.
Ware, Pierce, Glynn, Camden,
Charlton, Clinch, Echols, Lowndes (abundant)

{ Similar in habit to the preceding,
but rarely associated with it,
and less common. Ware, Glynn,
Charlton, Clinch, Lowndes.

{ Dry pine-barrens, etc.
Glynn: Vicinity of Dock Junction.
Camden: South of Woodbine.
Charlton: Sand-hills near Golleston.
Lowndes: Near Blanton.

Among wharves, Brunswick.

{ Dry or rather dry pine-barrens, common.
Ware, Glynn, Camden,
Charlton, Lowndes.

Charlton: Dry pine-barrens near
Camp Cornelia.

Chondrophora nudata (Hx) Britton.

Solidago

Euthamia caroliniana (L.) Greene.

Aster squarrosus Walt.

Leucocarpus bifolius (Walt.) Porter.

Erigeron vernus (L.) T. & G.

{ Moist pine-barrens, common
and often very abundant.
Ware, Pierce, Glynn, Camden,
Charlton, Blinch, Lowndes.

Crackish marshes of White Oak River.

{ A weed, along or near the O. & P. R. R.
Camden: West of Bull Head Bluff.
Charlton: Folkston, and westward.

{ Rather dry pine-barrens.
Ware: Near Waycross.
Camden: Woodbine; South of Kingsland
Charlton: Near Folkston & Camp Cornelia.

{ Dry pine-barrens, etc.
Camden: Between Woodbine & Colesburg
Charlton: Sand-hills S. W. of Folkston.

{ Moist pine-barrens, etc.
Ware: East of Waycross and Ruskin.
Camden: South of Colesburg
Blinch; Near Fargo.

Leptilon canadense (L.) Britton

" *linifolium* (Willd.)

Doellingeria reticulata (Pursh) Greene

Baccharis angustifolia Mx.

" *halimifolia* L.

A weed.

Glynn: Streets and wharves, Brunswick.

Camden: West of Bull Head Bluff.

Cumberland Island.

Charlton: Near Camp Cornelia.

Streets and wharves Brunswick.

Rather dry pine barrens.

Glynn: Near Cartier.

Camden: Woodbine; South of Kingsland.

Charlton: Near Holdston; East slope of ridge near Camp Cornelia.

Blinch: Near Fargo.

Brackish marshes of White Oak River.

Swamps, etc.

Glynn: Brunswick, and westward.

Camden: West of Bull Head Bluff;

Taxodium distichum swamps near Glencoe & Colesburg.

Along White Oak River.

Charlton: Along river at Traders Hill.

Swamp near Camden Co. line.

Not seen west of ridge.

Pluchea

" foetida (L.) C. & S.

" imbricata (Bearney) Nash

Pterocaulon undulatum (Walt.) Muhl.
(P. hysterochaeta (Mx.) Ell.)

Silphium compositum Mx.

Swamps, etc.

Glynn: Around artesian well, St. Simon's.
Between Bladen and S. B. T. crossing.

" Dock Junction and Carters.

Camden: Inner shore, Cumberland &
North of Colesburg.

Along Big Catfish Creek.

Not seen more than 25 miles inland.

Wet pine barrens and ponds.

Glynn: Near Dock Junction.

Camden: Frequent along railroad

Charlton: " west of "

{ Similar in habitat to the preceding
but never associated with it.
Ware, Camden, Charlton, Blinch.

Dry or rather dry pine-barrens.

Ware, Glynn, Camden,
Charlton (common), Blinch.

Camden: Dry pine-barrens south
of Kingsland.

Charlton: Sand-hills near Traders Hill.

Acanthospermum australe (L.) Moench.

Berlandiera tomentosa (Pursh) Nutt.

Rudbeckia hirta L.

" *Mohrii* Gray.
(*R. bupleuroides* Shutt.)

Borreria frutescens (L.) DC.

Helianthus

{ A common weed, especially along railroads.
Pierce: Fort Mudge and Hoboken.
Glynn: Among wharves, Brunswick.
Charlton: Racepoint, Camp Cornelia, Folkston.
Camden: South of Colesburg.
Lowndes: Valdosta.

{ Dry pine-barrens, etc.
Camden: Cumberland Island.
South of Kingsland.
Charlton: Sand hills near Folkston.

Camden: Dry pine-barrens
south of Kingsland.

{ Water-pine-barrens and ponds.
Ware: Between Bolen and Elsie.
Southwest and southeast of
Waycross.

{ Glynn: Salt marshes, Brunswick.
Camden: Inner shore, Cumberland I.

Camden: Moist pine-barrens
west of Bullhead Bluff.

Helianthus Radula (Pursh) T. & G.

Coreopsis

" *lanceolata* L.

" *nudata* Nutt.

"

Bidens

Baldwinia uniflora Nutt.

Actinospermum angustifolium (Pursh) Torr.

Rather dry pine-barrens, etc.

Ware: Near Waycross.

Camden: Frequent.

Charlton: Near Folkston.

Moist pine-barrens.

Ware: Between Waycross & Ruskin.

Camden: West of Bull Head Bluff.

Charlton: Northeast of Folkston.

Lowndes: Valdosta

Camden: West of Bull Head Bluff.

Ware: Cypress pond between
Waycross and Ruskin.

Charlton: Pine-barrens about
5 m. northeast of Folkston.

Satilla River swamp near Woodbine (1564)

Rather dry pine-barrens, common.
Seen in every county except
Pierce.

Pierce: Sand-hills of Satilla River.

Glomeria trinervia (Heng.) Mohr.

Marshallia graminifolia (Walt.) Small

Helenium tenuifolium Nutt.

"

Artemisia vulgaris L.

Erechtites hieracifolia (L.) Raf.

Carduus repandus (Mx.) Pers.

"

Thysanthera semiflorescens (Walt.) Runtz

THE END.

On ballast, Brunswick. (1521)

{ Moist or rather dry pine-barrens.
Ware, Pierce, Camden, Charlton.

A common weed, especially along roads and railroads.

Ware, Pierce, Glynn, Camden, Charlton, Echols, Lowndes.

Charlton: Pine-barrens about 5m. northeast of Folkston.


On ballast, Brunswick (1522)

{ A weed, often in swamps; from the islands to Okefinokee Swamp.
Glynn, Camden, Charlton.

Dry pine-barrens near Braganza.

{ Glynn: Streets of Brunswick
Camden: South of Woodbine.
Charlton: Along river at Traders Still.
Swamp near Camden Co. line.


Moist pine-barrens, not common.
Ware, Charlton, Clinch.



ITINERARY

of
Roland M Harper

August, 1902



JOURNAL

of a botanical exploration of
OKEFINOKEE SWAMP
and adjacent territory

in the counties of

Ware, Pierce, Charlton, Wayne,
Glynn, Camden, Clinch, Echols
and Lowndes in

GEORGIA

and portions of

Nassau and Baker in FLORIDA

during the month of

AUGUST, 1902

BY

ROLAND M. HARPER

1902.

August 1. Friday.

Arrived in Waycross on Atlantic & Birmingham RR. about 10:30 a.m. Changed cars before dinner.

Started for Satilla River at 1:20 p.m., following the A. F. & W. railroad the entire distance (3 miles).

Columbia formation covers practically the whole surface around Waycross, but Lafayette (and perhaps some underlying Tertiary) is exposed in several cuts toward the river.

Surface of the country normally very flat, but undulating between Waycross and the river.

A few specimens of Baptisia perfoliata seen on right-of-way about two miles from Waycross, but perhaps not indigenous here. Sagittaria Hobii also seen. Woods along here mostly composed of Pinus palustris and P. kubensis, with some P. serotina, but all considerably tampered with by the operations of

1 (continued).

turpentine, lumber, agriculture, etc.

Sand-hills of Satilla River (all on left side in Pierce Co.) are about a mile wide. At outer edge (i. e., farthest from the river) they are much like the ordinary sand-hills a little farther inland, being characterized by such species as

Cracca Virginiana

Pteridium

Quercus Catesbaei

Myrica humilis

Chrysobalanus oblongifolius

Serenoa serrulata

Botodendron arboreum

Opuntia vulgaris

Smilax humilis

Rhynchospora Grayi

Quercus cinerea

" *Virginiana*

" *Margaretta*

" " X *Virginiana* (?)

Rhus Toxicodendron

Eriogonum tomentosum etc.

I (continued)

In slightly moist spots the following species are met with:

Sophronantha hispida
Liospyros Virginiana
Rolisma ferruginea
Juncus sp. (no. 1445)

Toward the river the sand is whiter and drier, and covered with an almost impenetrable thicket of shrubs 6 to 10 feet high,

the following species predominating:

Polypodium stamineum ? (5-6 ft. tall)

Quercus sp. (no. 1466)

" *Virginiana*

Rhus copallina

Rolisma ferruginea

Osmanthus Americana

Batodendron arboreum

Hamamelis Virginiana

Castanea sp. (no. 1467)

Serenoa serrulata.

Very little herbaceous undergrowth in this thicket. Took ~~two~~ or three pictures of it.

I (continued)

Collected a Polanisia (no. 1464) and a Warea (no. 1465) on bare sand close to the railroad, but neither appears perfectly indigenous here. (The same is true of Ectinospermum angustifolium, which grows with them.)

Crossed the river again at 4 p.m. The Taxodium here seems to be all imbricarium. Nyssa ogeche is common here as on all the small rivers and creeks in this vicinity.

Collected 4 numbers and took 8 pictures on this trip. Got back to Waycross about 5 p.m. Rain commenced about 5:15 and continued rest of the evening.

Went to depot and met Ricker on Plant System train from Jacksonville at 5:25. Spent rest of day in discussing plans for exploration, etc.

2. Saturday.

Went for an all-day trip southwestward to Glenmore and beyond, starting from Waycross at 7:50 a.m. Sky partly cloudy, air very clear but not hot. Walked the railroad all the way, there being too much water to walk anywhere else.

Country almost perfectly flat, and only one curve on the railroad, about 4 miles from Waycross. (Track said to be straight for 30 or 40 miles beyond Glenmore).

Not much distinction between swamps, ponds and branches along here, and water pretty high everywhere.

Flora not very remarkable. Principal trees Pinus palustris, P. taeda, and Taxodium imbricarium.

Nyssa ogeche, Cephalanthus, Magnolia glauca, Nyssa biflora, etc. in creek swamps beyond Rushin. Pinus serotina quite abundant near Glenmore, forming an unmixed

2. (continued)

growth over small areas.

Rhynchospora corniculata is perhaps the most abundant herbaceous species in open swamps between Ruskin and Glenmore. Sarcocornia flava seen only once, near Ruskin. It seems to be very rare in this flat country (but grows in countless millions in the rolling pine-barrens a little farther inland). A good deal of Sabbatia decandra and some Rudbeckia Mohrii seen, in addition to the ordinary species of moist pine-barrens. Tasted rootstock of Dampiera for first time (taste indistinguishable), and helped Ricker collect some seeds of it.

Collected nos. 1468-1470a on this trip and took 3 pictures in the pine-barrens. No good water found along the route; wells all very shallow and warm.

No snakes or mosquitoes seen all day.

2. (continued)

Went about $1\frac{1}{2}$ miles beyond Glenmore (or 13 from Waycross), turned back about 4 p.m., and left Glenmore on train about 5, reaching Waycross at 5.30. Wrote up notes after supper.

3. Sunday.

Changed driers and put plants of two previous days in press. Wrote some letters and walked about the city some. Light showers during the day.

4. Monday.

Spent most of the morning buying supplies to use in the swamp.

Interviewed two men who had been in there hunting several times.

Changed driers and packed up.

Went out with Ricker about middle of afternoon to take some pictures.

4. (continued)

Left Waycross at 5:40 p.m., reaching
Folkston (33 miles) at 6:30.

Country very flat all the way, except
for a long cut 6-8 feet deep (in Columbia
sand) between Pacepond and Uptonville
and a few undulations along streams
near Folkston. Columbia surface
everywhere, no Lafayette seen.

Flora quite uniform all the way
from Waycross to Folkston. The
following are the principal species
seen from the train:—

Dichromena latifolia

Pontederia cordata

Eriocaulon decangulare

Taxodium imbricarium

Tillandsia usneoides

Serenoa serrulata (very abundant)

Pinus palustris

" *cubensis*

Xyssa biflora

Oxypholis filiformis

Chondrophora

Woodwardia Virginica

4. (continued).

Pterocaulon undulatum
Eupatorium rotundifolium
Cheria glabella
Eupatorium compositifolium
Pluchea sp. (no. 1430)
Magnolia glauca
Cheria Virginica ?
Cracca Virginiana
Quercus sp. (no. 1457)
Carduus repandus
Myrica humilis
Asclepias lanceolata
Sarracenia minor
Aristida stricta
Eupatorium album
Trilisa odoratissima
Galearia lanceolata
Juncus repens
" *trigonocarpus*
Castanea nana ?
Polycodium stamineum
Quercus Batesbaei
Pinus serotina (not common)
Vernonia angustifolia

4. (continued)

Elethra alnifolia
Polygala ramosa
Tabbata macrophylla
Lilium latesbaei
Hypericum fasciculatum.

Woods pretty dry most of the way,
 not full of water as near Alermore.
 Very few streams crossed.

Pines nearly all turpentine, also
 lumbered to some extent, but very
 little cleared land seen. ^{Country very} thinly settled.

Folkston seems to be about at the
 edge of the imperfectly developed sand-
 hills of the St. Mary's River, and
 the soil there is fine loose dry
 white sand. *Xelenium tenuifolium*
 and *Acanthospermum australe* common,
 especially the latter, which perhaps
 acts somewhat as a sand-binder.

Interviewed some of the natives about
 means of getting into Okefinokee Swamp.
 The swamp is described as a very
 interesting place, and this whole

4 (continued).

region is considered very salubrious, malaria being unknown. Dangers of the swamp have evidently been exaggerated by those who have never seen it. No one is known to have ever perished in there from snake-bite or any other cause.

5. Tuesday.

Water from pump at hotel strongly impregnated with iron, giving it a rather agreeable taste. Railroad water-tank supplied from an artesian well, the water from which rises to within a few feet of the surface.

Folkston has just been made county-seat of Charlton Co., and a new court-house and a few other new buildings are in process of erection.

5. (continued)

Spent the morning waiting for some of our baggage to come on from Waycross (it arrived about 11 a.m.) and in changing driers, etc. After dinner engaged a man to take us out to the swamp tomorrow morning, then walked the railroad down to St. Mary's River (3 miles away) and back. Topography undulating all the way, much like between Waycross and the Satilla River, but more sandy. Columbia formation 6 or 8 feet deep in many of the cuts, but not forming regular sand-hills as it does along the streams farther inland. Little if any Lafayette seen. Crossed river into Florida at 4:10 and back at 4:20. High bluff on Florida side, and railroad goes through it in a deep cut exposing Lafayette and older underlying strata. River here about the size and appearance of the

5. (continued)

Savilla near Waycross, but the Taxodium in it seems to be all T. distichum. Georgia side of the river here flat but not very swampy. In the river-bottoms about $\frac{1}{2}$ mile back is a rather peculiar association of plants, including the following species:

Scleria hirtella

Campulocarpus aromaticus

Koeleria hyssopifolia

Rhexia glabella (abundant)

Hypericum myrtifolium

Sabbatia campanulata

Ludwigia virgata (?)

Sarracenia minor

Sieglingia ambigua

Eryngium synchaetum

Crotonopsis spinosa (rather rare)

Rhynchospora Baldwinii

Dichromena latifolia

Quercus minima

Scleria gracilis

Most of these are plants characteristic of moist or rather dry pine-barrens.

14
5 (continued)

In damper places near by are found the following:

Taxodium imbricatum

Nyrica cerifera

Cladium affusum (all sterile)

Ludwigia linifolia

Pinus Sabensis

Clethra alnifolia

Asclepias lanceolata.

Got back to Holkston about 5.40.

Laid in some ^{more} food supplies (crackers & canned goods) to take into the swamp.

6. Wednesday.

Started for the swamp at 6.10 a.m., with J. W. Roddenberry and his two buggies and a driver. Obtained considerable information about common names and economic plants from Mr. Roddenberry during

6. (continued)

the drive. Two common weeds noticed in the outskirts of Goldston, Stenophyllus Floridanus called "water-grass" here as in Bullock and Coffee Cos.) and Erigeron Floridana ("poor-land weed"). Country very sandy and quite hilly the first few miles, becoming gradually flatter toward the swamp.

The following species are characteristic of the sand-hills near Goldston:—

Eriogonum argyranthemus

Psoralea canescens

Quercus Catesbaei ("black-jack")

" *Margaretta*

Eriogonum tomentosum

Chrysobalanus oblongifolius

Quercus cinerea ("turkey oak")

Asimina sp. (no. 1438)

Breueria humistrata

Quercus minima

Jatropha stimulosa

Cracca Virginiana ("devil's shoestring")

(The juice of the root said to poison fish)

6 (continued)

Polygala grandiflora

In moister places near one of the first streams crossed are seen:

Quercus digitata ("red oak")

Diospyros Virginiana

Liriodendron Tulipifera

Pinus serotina

Rhus Vernix

Cephalanthus

Cynilla

Magnolia glauca

Alnus incana

Syalea viscosa

Woodwardia melata

Phoradendron flavescens

Gillilandia asneoides

Serenoa serrulata

Ilex glabra

Smilax laurifolia

Flora quite rich all the way, and several unfamiliar plants seen. Pine forests pretty badly

6 (continued)

umbered. Road becomes rather indefinite last few miles, and has apparently not been traveled for several days. No shoes on horse, not needed in this pure Columbia sand. Houses pretty scarce. For the last three or four miles road goes over a broad smooth sandy scarcely perceptible ridge bounding the eastern margin of the swamp. The following species are common on this ridge: -

Sarracenia minor
 " *psittacina*
Serenoa serrulata (very abundant)
Centella Asiatica
Rhexia glabella
Galactia Elliottii
Oxypolis filiformis
Rhexia villosa
Lilium Batesbaei
Doellingenia reticulata
Xyris torta
Azilia cassioides

6 (continued)

Chrysolanus oblongifolius
Polygala setacea
Himbiistylis puberula
Lapodium imbricarium (in depression)
Cyperus filiculmis
Peridium
Aristida stricta
Ergonium aromaticum
Asclepias cinerea
Elephantopus scaber
Pinus palustris (fruiting abundantly)
Myrica parvifolia
Ralmia hirsuta
Xyris brevifolia
Xolisma ferruginea

Reached Camp Cornelias, on the canal, at the edge of the swamp, at 9.35 a.m. One family still lives there, to look after the property belonging to the owners of the swamp.

Started out a few minutes later to find a guide living about

6 (continued)

three miles north of Camp Cornelia, and engaged him for \$1.50 a day. Borrowed a machete and a tin fly from the office, and started down the plank walk (made of Taxodium imbricarium) to the swamp at 2:25. Passed several empty houses and ruins of sawmill and terminus of extinct Brunswick & Pensacola R.R. (which formerly extended from here to Bull Head Bluff on the Satilla River, passing through Folkston).

Embarked with Ricker and the guide in a small boat on the canal at 2:55. Had our pictures taken with smallest camera just before starting.

Canal at this point full of wrecks of steamboats, dredges, etc. Canal said to have been originally 40 feet wide and 15 feet deep, but now somewhat smaller on account of caving in of banks and

(continued)

growth of aquatic vegetation, such as Pontederia, Sagittaria, Crontium, Eleocharis elongata, etc.

Swamp very open, for first few miles at least. Few trees seen except Taxodium imbricarium, which gives little shade. Pinus kubensis occasionally seen. Tillandsia usneoides always in sight. Banks of canal lined with Panicum digitarioides all along, accompanied by Crontium, Pontederia, Gyneroschea, etc.

Banks variable in height, but usually too high to see over while sitting in boat, principally on account of dense growth of Panicum digitarioides.

At several points not far from entrance of canal are remains of experimental gardens on the elevated banks, but nothing now growing in them but Panicum digitarioides. Soil seems to be mostly peat, and banks of canal were seen on fire in two or three places.

At several places along the canal are numerous muscadine vines (Vitis rotundifolia)

6 (continued)

var.?), bearing abundant ripe fruit at this time. These are said to have all come from scuppernong seeds thrown out by the crews of the various boats which formerly navigated the canal. Fruit is dark purple, but smaller than either the cultivated scuppernong or the native muscadine, and ripens unusually early.

Sunny day, but cool and pleasant. Only one snake seen, and no mosquitoes until nightfall. Reached Bugaboo Landing, a slightly elevated sandy place on canal bank, about 7 miles from Camp Cornelia, at 6:10 p.m., and camped there for the night. Got water out of a sphagnum bog just over the bank. No discomforts worth mentioning yet.

7. Thursday.

Night passed off without incident, but mosquitoes rather troublesome. Got up about 4:30 a.m. Collected nos.

7 (continued)

1471-1473a in the immediate vicinity,
and started on foot for Bugaboo Island
(about 2 miles west or southwest) at 6:50.

Most of the way is through sphagnum
bogs (into which we often sank knee-
deep), becoming gradually drier toward
the island. Herbaceous vegetation of these
bogs consists mostly of:-

Woodwardia Virginica (very abundant)

Drosera sp. (no. 1472)

Eriophorum Virginicum ("Indian cotton")

Dulichium arundinaceum

Sarracenia minor (leaves 2-3 ft. tall)

" *pittarina* (also very large)

Limnophyton tuberosum

Peltandra sagittifolia

" *Virginica* (rare)

Rhynchospora sp. (no. 1477)

Of shrubs and vines the following species
prevail:-

Pieris nitida

" *phillyreaefolia*

Persea

Smilax laurifolia

7 (continued)

Ilex glabra
Itea Virginia
Leucothoe elongata?

The only trees noticed here are *Pinus*
Bubensis (abundant but rather small)
and *Taxodium imbricarium* (less common)
Reached island about 8 a. m. Boundary
between it and the surrounding bog
very indefinite and scarcely perceptible,
the island being simply an area where
the sand reaches the surface. It is
said to be about two miles in diameter,
and was discovered only a few years
ago. Flora not rich, consisting of
the following species:—

Pinus Bubensis (everywhere)

" *serotina* (rather rare)

Serenoa semulata (everywhere on island,
not seen elsewhere in swamp)

Gaylussacia frondosa

Kalmia hirsuta

Pieris nitida

Vaccinium nitidum

Quercus minima

7 (continued)

Zolisma ferruginea (only knee high)*Ilex lucida*" *glabra**Laylussacia dumosa**Chrysopsis graminifolia**Galactia Elliottii* (rare)

Took several pictures on the island, but didn't collect anything there. Collected nos. 1474-1476 on way out and nos. 1476a-1478 on way back by same route. Reached landing at 10:30 and re-embarked on canal at 11 a.m.

Saw *Pieris phillyreaefolia* climbing *Taxodium* to a height of 20 feet or so by creeping under bark. Collected specimens (no. 1479) and photographed it.

Came to edge of Chase Prairie about noon. The prairie is an open area several square miles in extent, mostly covered with shallow water, and dotted with clumps of *Taxodium* (called "houses"). No *Sphagnum* in prairie, but ground

7 (continued)

very boggy. Following species noted:-

- Eriocaulon compressum*
- Scleria* sp. (no. 1480)
- Crontium aquaticum*
- Rhynchospora corniculata*
- Triadenum Virginicum*
- Drosera* sp.
- Peltandra Virginia*
- Daphnys flavidula*
- Sarracenia minor* (very large)
- Panicum digitarioides*
- Castalia* sp.
- Gyneroschea tinctoria*
- Pontederia cordata*
- Sagittaria* spp.
- Utricularia purpurea*
- " sp. (no. 1481)
- Limnanthemum aquaticum*

Camped for the night in a small shack in the edge of Jackson Bay, just beyond Chase Prairie, and about 11 miles from Camp Cornelia. Rested more comfortably than last night. Mosquitoes fewer.

8. Friday.

Got up about 4:30 a.m., and collected nos. 1481 b-1482 v in the bay before breakfast. Bay consists of a dense forest of *Taxodium imbricarium*, with the following shrubs and herbs:-

Ilex cassine

Rhus radicans

Cynilla racemiflora

Tillandsia usneoides

Xyris fimbriata

Smilax laurifolia

Pieris phillyreaefolia

Woodwardia Virginica

Itea "

Panicum digitarioides

Sarracenia minor

Peltandra Virginica

Pieris nitida

Eriocaulon compressum

Persea

Magnolia glauca

Left camp at 6:05 a.m. and started for civilization again. Collected

27
8. Friday

nos. 1483 and 1484 along the canal on the way out. Disembarked at 11:50 and reached the settlements at Camp Cornelia a few minutes later.

Afternoon cloudy. Went south of Camp Cornelia about $\frac{1}{2}$ mile. Country there very flat, with a few shallow cypress ponds. Many fine young saplings of Pinus palustris seen near the house. Stenophyllus stenophyllus abundant in unfrequented road. Collected nos. 1485-1489.

After supper found a number of discarded letter-copying books containing correspondence of the defunct Suwannee Canal Co., and made copious selections therefrom. Saw also some maps (both plan and profile) of Brunswick & Pensacola RR., but data on them rather meager.

Noticed a large specimen of Sambucus canadensis (still in flower) in front yard. Said to be small in comparison with some specimens in creek swamps.

9. Saturday.

Heavy rain at daylight, continuing without interruption until about 1 p. m. Made further selections from Swane's Canal Co.'s correspondence.

Started out with umbrella about 10:45 and went two or three miles north through extensive pine-barrens in the rain. Collected nos. 1490 and 1491. Got back about 1:15 p. m.

Took some 4x5 pictures in the vicinity of Camp Cornelia. Started alone for Folkston at 4:15 p. m., following ~~extinct~~ B. & P. R. R. most of the way, carrying nothing but portfolio. Left Ricker and the baggage at Camp Cornelia to follow later. Track straight for about six miles. Ridge bounding the swamp scarcely perceptible in walking over it, but quite conspicuous when looking back along the track from a few miles to the eastward.

Pine forests badly lumbered most of the way, but many young trees now springing up.

21
9 (continued)

Collected no. 1412. Turned off to left on dirt road near Trader's Bluff about 7 p.m. Pretty good sand-hills along here, but too dark to study their flora. Reached Holston at 7:55.

10. Sunday.

Ricker arrived with baggage at 7:45 a.m. Changed all driers and put Chefinokee plants in press. Spent several hours writing up notes, etc.

11. Monday.

Ricker left for St. Augustine at 7:40 a.m. Started for Trader's Bluff at 8, by the road. Sand-hills nearly all the way, except first $\frac{1}{2}$ mile and vicinity of Spanish Creek and one of its tributaries, where there is a flora much like

11 (continued)

that of St. Mary's River bottoms.
 Reached Trader's Hill about 11:15.
 It is a small, old, settlement in a
 grove of Quercus Virginiana on a
 bluff of the St. Mary's River 50
 feet high or more. Went down to
 river and crossed it on trestle of
 extinct St. Mary's, Lake City & Gulf RR.
 Several interesting plants seen along
 river. Took several 4 x 5 pictures.
 Started back about 1 p. m. Noticed
 the following species on sand-hills:-

Eriogonum tomentosum
Chrysobalanus oblongifolius
Ruhmisteria pinnata
Quercus batesbaei
Polygala grandiflora
Cracca Virginiana
Quercus Margaretta
Baptisia lanceolata
Croton argyranthemus
Breweria humistrata
Serenoa serrulata

11 (continued)

Quercus cinerea
Polycolium stamineum
Cassipouma pectinata
Asimina sp. (no. 1438)
Chamaecrista fascicularis?
Sericocarpus bifolius
Carphephorus sp.
Eryngium aromaticum (no. 1496)
Juncus biflorus (no. 1495)
Psoralea canescens
Helianthus Radula
Meibomia arenicola
Scleria sp. (no. 807)
Jatropha stimulosa
Cyperus filiculmis
Stylosanthes biflora
Asclepias cinerea
Vernonia angustifolia
Berlandiera tomentosa

3

In the peculiar flats near Spanish Creek the following species were noticed:—

11 (continued)

- Taxodium imbricarium* (scattered)
- Rhexia glabella* (the most characteristic and abundant species in this "formation")
- Rhexia lutea*
- Thyphocarpum myrtifolium*
- Eriocaulon decangulare*
- Sarracenia minor*
- Sabbatia campanulata*
- Scleria gracilis*
- Rhynchospora semiplumosa*
- Campulocarpus aromaticus*
- Dichromena latifolia*
- Sieglingia ambigua*
- Scleria hirtella*
- Pinus cubensis*
- Chondropora* (not common)
- Koelia hyssopifolia*
- Polygala ramosa*
- Panicum stenodes*

(Compare with list on p. 13 of this book).
 Followed B. & O. R. R. most of the way
 back to Holston (3 or 4 miles).
 Grade descends rapidly toward

11 (continued)

Spanish Creek, and rises rapidly on other side. Considerable Lafayette exposed beneath Columbia sands in cuts on both sides of creek. Collected nos. 1493-1505 on this trip. Got back to Holston at 3:15.

12. Tuesday

Started at 6:33 a. m. for Bull Head Bluff on Satilla River in Camden Co., said to be about 10 miles away. Followed extinct B. & P. R. R. all the way except first half mile. Country all rather dry pine-barrens and perfectly flat for two miles or so; then small sand-hills and one or two branches, with sphagnum bogs along first one, containing the following species:-

Oxocera sp.

Ilabenaria integra (no. 1507)

Eriocaulon decangulare

Tofieldia racemosa

Burmannia capitata

12 (continued)

Marshallia graminifolia
Campulocaulis aromaticus
Saururus cernuus
Xyris Baldwiniana
Thysanotus semiflosculus
Lycopodium Carolinianum
Utricularia juncea
Chondrophora
Asclepias lanceolata
Polygala lutea
Centella asiatica
Melanthium sp. (no. 1506)
Mayaca subletii
Habenaria blephariglossa
Limnium tuberosum
Bartonia virginica

In wet woods between the bog and the branch
are the following: -

Magnolia glauca
Rhus Vernix
Ilex lucida
Azolla racemiflora
Najas Carolinensis
Samolus regalis
Woodwardia Virginica

12 (continued)

Second branch runs through a swamp having quite a different flora, such as:

Nyssa biflora
Phoradendron flavescens (on the preceding)
Alnus rugosa
Smilax laurifolia
Acer rubrum
Tillandsia sp. (no. 1509)
Polypodium polypodioides
Rhus radicans (got poisoned by it)
Crontium aquaticum
Euscuta compacta

Photographed the *Tillandsia* before collecting it. Country more undulating beyond these branches, but track nearly straight all the way. *Taxodium imbricarium* not seen until 9:40, 3 or 4 miles from Holston. *Pinus serotina* frequent along here. Made some photographs of it. Beyond this the railroad passes through several swamps of various sizes, the flora of which seems to indicate absence of Lafayette formation. No ponds seen

12 (continued)

all day. Cladium effusum seen at frequent intervals, but nearly always sterile except in ditches or other places where ground had been disturbed. Crossed a tram-road in operation about 5 miles out. Several unfamiliar plants seen just beyond this point. About 7 miles from Folkston, just west of Camden Co. line, is a particularly interesting swamp about $\frac{1}{2}$ mile wide, with no water flowing through it at this season. Flora composed of many species of deciduous trees, and varied herbaceous vegetation. Path through it nearly choked up with weeds and briars. Came out of swamp about 12:10 and saw house and field ahead (first seen for several hours). Reached house at 12:30 and got some water (first obtainable since leaving Folkston) from the well. Well is warm, muddy and shallow, but said by owner (a colored man) to be inexhaustible owing to a fissure (probably in the Lafayette

12 (continued)

clay, 2 or 3 feet below the surface) caused by earthquake of 1886.

Reached Bull Head Bluff at 1:30.

Bank of river at this point, about 8 feet high, with exposures apparently of Columbia, Lafayette, and ^{a little} underlying Tertiary. Saxodium distichum grows here at edge of water just below the Lafayette, and S. imbricarium about 200 yards back from river, doubtless on the Lafayette.

Two or three houses and a steamboat landing on this side of river. Opposite bank low and swampy, lined with large specimens of Pontederia cordata, and other plants unrecognizable at this distance. There is some tide at this point, and steamboat is said to pass about 3 times a week.

Left Bull Head Bluff at 2:30 and went back to Holkston by same route. Pine forests pretty badly lumbered most of the way, and somewhat turpentine, though many

12 (continued)

square miles uninhabited.

Collected nos. 1506-1517 on this trip and took six $2\frac{1}{4} \times 3\frac{1}{4}$ pictures and 26 pp. of notes. Got back to Folkston at 6.50.

13. Wednesday.

Put plants in press, changed driers, and packed up. Found that local train on Jesup Short Line only runs Tuesday, Thursday and Saturday, so had to go around by Waycross to get to Brunswick.

Left Folkston at 4:20 p. m. and Waycross at 5:40. Country east of Waycross pretty flat. Sand-hills along a few of the streams. Plenty of Taxodium imbricarium seen. Eriogonum tomentosum last seen in Pierce Co., near Globoken, 44 miles from Brunswick. No Parracenia flava seen for a week or more. Crossed Jesup Short Line at Nahunta, 6:23 p. m.

13 (continued)

Its track is perfectly straight beyond horizon in both directions, showing level nature of country. Crossed Satilla River at 6:41. Extensive swamps on both sides of it here, apparently no sand-hills. Too dark to see much beyond this point. Passed Bladen at 6:57 and saw Sabal Palmetto there. It is frequent rest of way to Brunswick. Plenty of salt marshes near Brunswick. No sawmills or turpentine stills seen on this road east of Waycross. Reached Brunswick about 7:30 p. m.

14. Thursday.

Walked about the city from about 6 to 10 a. m., noting the weeds in the streets, etc. Went also about 2 miles back up the railroads, and into salt marshes east of city. Flora of salt marshes much like those of Chatham Co.

14 (continued)

Principal business at Brunswick seems to be exportation of the products of Pinus palustris (and P. bubensis), especially cross-ties.

15. Friday.

Went down railroad to lower end of city, about 2 miles beyond depot. Collected nos. 1518-1527 (mostly ballast plants) along the rosin wharves. Came back on other side of city and collected nos. 1528 and 1529. Two or three fine flowing artesian wells along the route.

Ricker arrived on Brunswick & Birmingham R. R. at 12:46. Oiers shipped from Holston not yet arrived. Not much sunshine all day. Thunderstorm about 6 p. m.

16. Saturday.

Put yesterday's plants in press.
Embarked for St. Simons Island at
8:35 a. m., landing at Ocean Pier,
at south end of island, at 9:20.
Walked from there east and north
along the beach about 2 miles, to
an inlet too wide to cross, and
back by ~~same~~ route. Several
fine artesian wells on the island.
Serenoa abundant near Ocean Pier
(probably the same place where William
Bartram saw it in 1774), but
not seen elsewhere on island.

No Sabal Palmetto seen, and man-
^{other} characteristic plants of Tybee Island
apparently absent.

Cottages on the island mostly empty
and many of them twisted around by
cyclones and otherwise dilapidated.
Several cultivated fields and permanent
habitations seen.

Salt marshes present, but not
much variety in their flora.
Saw beds of oysters on stiff clay

16 (continued)

exposed at low tide. No mosquitoes seen all day.

Collected nos. 1530-1535 and took 7 pictures on the island. Plant communities seen may be roughly classified as follows:-

Beach plants.

Euphorbia polygonifolia

Polygonum maritimum (no. 1534)

Dune plants.

Portulaca maritima

Sesuvium serrulata

Suaeda imbricata

Salsola Kali

Yucca aloifolia

Xanthoxylum clava-herculis

Juniperus sp. (no. 1532)

Chenopodium sp. (no. 1535)

Thicket back of dunes

Callisarpa americana

Myrica cerifera

Suaeda frutescens

16 (continued)

Juniperus sp. (no. 1532)
Smilax sp. (no. 1531)
Paronychia sp. (no. 1533)
Quercus Virginiana
Bumelia Canadensis
Arenaria "
Opuntia Pes-Conv. (?)
Oplismenus setarius
Melothria pendula
Ampelopsis arborea

Salt marshes

Spartina maritima
Salicornia ambigua
Suaeda sp.

In and around pool formed by water from artesian well.

Chara sp. (no. 1530a)
Hydrocotyle umbellata
Monnina Monnina
Paspalum distichum
Cyperus flavescens
Lippia sp.

16 (continued)

Pluchea camphorata
Mikania scandens
Labbatia stellata

Ruderal plants

Argemone alba (no. 1530)
Cassia Tora
Erechtites hieracifolia
Capriola Dactylon
Ambrosia artemisiifolia

Left the island at 5:40 p. m. and
reached Brunswick at 6:30. Heavy
rain and wind on return trip.

17. Sunday.

Too cloudy to to change driers.
Assorted some specimens already
dried. Wrote some letters.
Went down on wharves, etc., with
Ricker.

18. Monday.

Went to custom-house to look at some charts of the coast.

Started out at 2:10 p.m. and went up the railroad (Plant System & Southern combined) beyond Dock Junction (about 7 miles). Flat pine-barrens much like those in Charlton Co. begin about 3 miles out, and Lafayette formation probably reaches its seaward limit at this point. Plenty of Pinus cubensis and many small shrubs (especially Xolisma ferruginea) seen here. Taxodium imbricarium found in a shallow depression 4 or 5 miles from Brunswick, near Dock Junction. Saw some puzzling pines along here, one of them probably the true P. heterophylla of Elliott. Collected some specimens of it (no. 1537) about sunset, with great difficulty, also no. 1536 and 2 fungi. Got back to Brunswick at 7:50 p.m.

19. Tuesday

Wrote to Washington for some Coast Survey charts of the Georgia coast.

Embarked for Cumberland Island at 8:06 a. m. Looked through Ricker's field glasses from deck of boat at woods and marshes south and east of Brunswick. Saw some magnificent specimens of Sabal Palmetto along edge of woods and in separate groves, two or three miles away across the marshes. (This species is rather rare around Brunswick, and was not seen on either of the islands visited.)

Boat stopped at Jekyll Island at 9:50. No one allowed to land here without special permission. Several fine buildings near landing, and extensive forests all over the island. Plenty of Uniola paniculata seen on south end of Jekyll and north end of Cumberland, but none accessible to us.

Landed at High Point, on western shore of Cumberland, at 10:30.

19 (continued)

Edge of island here is a steep bluff about 25 feet high. On top of this bluff, and extending back $\frac{1}{2}$ mile or more, are dry pine-barrens similar to those of Charlton Co., with the following flora:-

Pinus palustris
Quercus Virginiana
Galium hispidulum
Galactia Elliottii
Juniperus sp.
Polypodium polypodioides
Tillandsia usneoides
Euphorbia cordifolia
Asimina angustifolia
Vernonia "
Elephantopus elatus
Bumelia lanuginosa
Gornia bracteata
Jatropha stimulosa
Clitoria Mariana
Quercus batesbaei
Psoralea canescens
Houstonia rotundifolia

11 (continued)

Eupatorium compositifolium
Perlandiera tomentosa
Ascyrum sp.
Stillingia sylvatica
Breweria humistrata
Stylosanthes biflora
Serenia serrulata (not common)
Rhynchospora Grayii (no. 1538)
Polygonella sp.
Sporobolus juncus
Morongia sp.
Simina sp. (no. 1438)

At edge of these pine-bayous is an abundance of *Bejania racemosa* (reported from this island by Elliott). Collected and photographed it. Just beyond is a sort of scrub thicket $\frac{1}{2}$ or $\frac{3}{4}$ mile wide, with flora much like that of the Satilla River sand hills in Pierce Co., but less xerophytic. The following species were noted here:

19 (continued)

Quercus sp. (nos. 1466, 1541)*Xolisma ferruginea* (10 ft. tall)*Pieris nitida*? (no. 1541)*Bejania racemosa**Serenoa seculata**Myrica cerifera**Pinus Buhensis**Dicranum* sp. (no. 1541a)

Followed mule-car track (narrow-gauge) most of the way across the island, a distance of about 2 miles.

Reached the beach about 1:30 p. m. Nobody in sight. Fine flowing artesian well on the beach, water from which is absorbed into the hot sand before reaching the ocean.

Wind pretty strong here, blowing constantly toward the land. Its effects are strikingly shown in the leaning of the stunted trees, and movement of the dunes. Dunes are now about $\frac{1}{2}$ mile from beach and advancing rapidly inland.

19 (continued)

Their encroachment on the forest behind them can be plainly seen. Between the beach and present line of dunes are remains of old dunes, capped and held in place by Leavenworthia serrulata.

On the shifting sands near the beach the following species prevail:-

Fimbristylis sp. (no. 1544)

Spartina juncea (no. 1545)

Craton maritimus

In the meadow a little farther back are seen the following:-

Dichromena colorata

Juncus megacephalus

Cyperus cylindrostachys

Sabbatia stellaris

Ludwigia sp. (no. 1542)

Gerardia sp. (no. 1543)

Oenothera humifusa

Scirpus Americanus

Juncus sp. (no. 1546)

Cyperus sp. (no. 1547)

Lippia sp.

17 (continued)

Sesuvium Portulacastrum occurs in moist hollows, which are evidently remains of pools which the dunes have passed over and left behind, partly buried. On the dunes the following species were noticed:-

Sesuvium serrulata

Eroton maritimus

Xanthoxylum clava herculis

Fimbristylis sp. (no. 1549)

Leptilon canadense (introduced?)

Salsola Kali

Euphorbia polygonifolia

Vitis rotundifolia

(This seems to act as a sand-binder)

Dunes quite broad and perfectly bare in some places.

Went from hotel down to landing on mule-car. Explored inner shore of island at base of bluff while waiting for boat, and noticed the following species:-

19 (continued)

Scirpus Americanus
Eroton maritimus
Pluchea camphorata
Monnina Monnina
Lippia sp.
Hydrocotyle umbellata
Cyperus sp. (no. 1547)
Juncus Roemerianus
Erechtites hieracifolia
Eleocharis sp. (no. 1550)
Distichlis spicata
Dondia sp.
Borrichia frutescens
Salsola Kali
Polypremum procumbens
Leptilon Canadense.

Collected nos. 1538-1550 and took
 16 pictures (with both cameras) on this trip.
 Embarked at 5:23 p. m. and
 reached Brunswick at 7:25.

20. Wednesday.

Orders shipped from Holston a week ago not arrived yet. Had agent telegraph for them.

Ricker left on Brunswick & Birmingham RR. at 10:40 a. m.

Changed all orders and put most of yesterday's plants in press. Evening unusually hot and sultry.

21. Thursday.

Packed up, etc. all morning, and just caught Brunswick & Birmingham train at 3:30 p. m.

Flat pine-barrens begin within a mile or two, and extend with few interruptions all the way to Thalmann (18 miles) with few interruptions.

Sabal Palmetto nearly always in sight, sometimes in dense woods and sometimes in open marshes.

Serenoa abundant all through the pine-barrens. Taxodium

21 (continued)

imbricaria scattered all along.
Pinus serotina seen at many points.
 Passed several interesting salt marshes
 and extensive swamps, containing
 unfamiliar plants. Reached Thalmann
 at 4:28 and left on F. C. & P. Ry.
 (now Seaboard Air Line) at 4:43.

Beautiful specimens of Sabal Palmetto
 abundant in the pine-forests around
 Thalmann and several miles south
 of there. Trunks rather slender,
 often 30 feet tall or more, usually
 naked. Many specimens in fruit.
 Plenty of "round timber" (mostly Pinus
palustris) between Bladen and S. B. F.
 crossings. In Camden Co. noticed
 numerous extensive swamps similar to
 those between Folkston and Bull Head
 Bluff, containing Taxodium distichum
 and other species presumably indicating
 absence of Lafayette formation.
 Reached White Oak about 5:20
 and stopped there a day.

Started at 5:30 for White Oak River,

21 (continued)

a small tidal stream, bordered by extensive brackish marshes, crossing the railroad just north of station. Tide rises and falls about 6 feet here, and river is navigable for sailboats up to railroad trestle. Collected nos. 1551-1553 in these marshes, and returned at 6:30. Water in river said to be alternately fresh and salt. Few mosquitoes seen here.

White Oak has fine artesian well system for such a small place (about 100 inhabitants), and has quite a thrifty appearance.

22. Friday.

Changed all driers and put yesterday's plants in press.

Saw a lot of Melochia corchorifolia in back yard. This weed is said to have come in about two years

22 (continued)

ago, and flourishes in cultivated land.
Other weeds noticed here are:

Cyperus *tria*
" *compressus*

Portulaca *oleracea*

" *pilosa*

Hollugo *verticillata*

Eleusine *Indica*

Euphorbia *maculata*

Commelina sp.

Saw ^{inverted} cypress knee (probably *T. distichum*)
used for trough under faucet in back
yard.

Started for White Oak River again
at 2:30. Collected nos. 1554-1558
and took three 4x5 pictures. The following
are representative species in the marshes:-

Eleocharis sp. (no. 1550)

Nipania *scandens*

Dichromena *colorata*

Sagittaria *lanceifolia*

Gizania *aquatica* (Panicle 4-5 ft.
long. Stems 10 ft. tall. Leaves about 7 ft. x 3 in.)

22 (continued)

Pontederia cordata (5-6 ft. tall)*Cyperus flasper*

" sp. (no. 1552)

" *virens**Spartina polystachya* (10 ft. tall)*Ludwigia alata* (no. 1554)*Cicuta Burtissii**Solidago sempervirens* ?*Lythrum lineare* (no. 1553)*Cladium effusum* (8 ft. tall)*Juncus Roemerianus* (6 " ")*Samolus floribundus**Kosteletzkya Virginica**Scirpus* sp. (no. 1556)*Scirpus* sp. (no. 1557)*Typha latifolia**Baccharis halimifolia*" *angustifolia*

Got back to depot about 5 p. m. and

left on southbound train at 5:17.

Extensive swamps again between White
Oak and Woodbine, as well as plenty
of flat pine-barrens. Marshes of

22 (continued)

Satilla River much narrower than those of White Oak River, but river considerably larger, with drawbridge across it.

Reached Woodbine ($4\frac{1}{2}$ miles from White Oak, and $\frac{1}{2}$ mile south of Satilla River) at 5:25. Seems to be a little smaller place than White Oak. Same kind of artesian water here as throughout the maritime counties, slightly sulphurous. Thunderstorm after supper.

23. Saturday.

Started out at 7:15 a. m. and went four miles down railroad and back. Track straight all the way and beyond horizon southward. Country nearly flat with very slight undulations, shallow ponds and sluggish streams. Pine forests lumbered and turpented a little, but almost unbroken. Pinus palustris and P. cubensis both abundant, and Taxodium imbricarium frequent.

23 (continued)

Lafayette formation exposed in a few cuts, under about 3 feet of Columbia sand.

The following species are characteristic of these flat pine-barrens:-

- Marshallia graminifolia*
- Triplaris odoratissima*
- Hypericum myrtifolium*
- Centella asiatica*
- Elocharis Baldwinii*
- Aristida spiciformis*
- Eupatorium rotundifolium*
- " *hyssopifolium*
- Ilex glabra*
- Serenoa serrulata*
- Xolisma ferruginea*
- Doellingeria reticulata*
- Bejania racemosa* (rather rare)
- Salmia hirsuta*
- Pterocaulon undulatum*
- Hypericum fasciculatum*
- Rhexia ciliosa*
- " *glabella*
- Baldwinia uniflora*

23 continued)

Galactia Elliottii
Sabbatia "
Sarracenia minor
Campulocarpus aromaticus
Cladium effusum (sterile)
Zabernia blepharigottis
Mesosphaerum rugosum
Polygala lutea
Eriocaulon decangulare
Pinus serotina
Ayris torta
Eupatorium verbenaeifolium
Hypericum opacum
Oxypholis filiformis
Oldenlandia
Lilium batesbaei
Coreopsis angustifolia ?
 etc.

Passed Colesburg (3 miles from Woodbine) at
 9:57 and again at 10:55. No settlement
 there except a store (including postoffice)
 and a house. Plenty of "round timber"
 (*P. palustris*) in the vicinity.

Collected nos. 1559 and 1560 on

23 (continued)

this trip. Got back to Woodbine at noon.

Went out between 1 and 2 p.m. and took 3 pictures along river with 4x5 camera. Started out again at 2:05 with portfolio and small camera. Took 4 pictures. Borrowed a boat and collected a *Spartina* (no. 1561) in edge of water. *Scirpus Lacustris* (about 8 ft. tall) grows with it; not seen elsewhere in Camden Co.

Marsh is bordered in some places by a belt of *Isenardia palustris* about 10 feet wide, succeeded by *Pontederia*, *Zizania*, *Sagittaria latifolia*, etc.

Took refuge in bridge-tender's house during a rainstorm from 3:30 to 4:45. Met there a gentleman who wants list of plants observed in Camden Co.

Went across river after rain and collected nos. 1562-1565 in the swamp just beyond the

23(continued)

brackish marsh. Flora here differs somewhat from that of the marshes, consisting (in part) of the following species:

Polygonum arifolium (no. 1563)

Eupatorium serotinum

Sambucus Canadensis

Typha latifolia

Comandra virginica

Mikania scandens

Ludwigia alata

Sidens sp. (no. 1562)

Juncus sp. (no. 1564)

Peltandra virginica

Scirpus eriophorum

Rhynchospora miliacea

Baccharis halimifolia

Panicum sp.

Magnolia glauca

Returned to Woodbine at 6:15 p. m.

24. Sunday.

Pretty cloudy morning. Started out at 8 a.m. and went west of Woodbine about a mile, discovering there a fine rice plantation on the right bank of the Satilla River, on a bluff about 10 feet high. Rice fields in marshes on opposite side of river, which are about $\frac{1}{2}$ mile wide. Going to a big bend in the river at this point, marshes are all on one side.

Beautiful view up and down the river at this point. Bluff presents about the same geological section as at Bull Head Bluff, which is on same side of river about 10 or 15 miles above. Taxodium distichum grows at edge of water at both places. Country between here and Woodbine much like that near Camp Cornelia in some respects.

Returned to Woodbine about 9:30, and went out about $\frac{1}{4}$ mile northeast of the station shortly before noon. Saw a single specimen

24 (continued)

of Podostigma in pine-barrens there. Changed all diuers after dinner (hindered considerably by wind and clouds), and put yesterday's plants in press. Cool and pleasant about sunset.

25. Monday.

Left Woodbine on southbound train at 7:50 a. m. Got off at Kingland (about 12 miles) at 8:09. Left most of hand-baggage in depot and started immediately for St. Mary's River, 4 miles down the railroad. Dense pine-barrens all around Kingland. Several small Sagittaria biflora ponds south of there.

A mile or two south of Kingland are two interesting creeks, Little and Big Catfish, the latter a tidal stream, but with only a narrow fringe of brackish marsh along it. Raxodium distichum occurs there. Between these creeks is a fine geological section

25 (continued)

of the two superficial formations, where railroad passes through a low sandy ridge in the pine barrens. Flora of this ridge the most xerophytic seen on the mainland of Camden County.

Came to edge of river swamp (which is about a mile wide here) about 10 a. m. Swamp here is different from all other river swamps seen, consisting mostly of a very open forest of *Pinus Cubensis*, with a dense undergrowth of the following woody plants:—

Gordonia Lasianthus (abundant)

Pieris nitida

Smilax laurifolia

Persea sp.

Ilex glabra

and the following herbs:—

Woodwardia Virginica

Sarracenia minor

Gyneria tinctoria

Rhynchospora sp. (no. 1477).

25 (continued)

At least four strata of vegetation are plainly discernible here, viz., the pines, the shrubs (all about 6 feet tall, Gordonia predominating), the herbs, and the sphagnum which covers the ground everywhere.

The whole has an aspect very similar to that of the bogs in the vicinity of Bugaboo Island in Okefinokee Swamp. (Compare with lists on pp. 22 and 23 of this book).

Toward the river the woods become denser and the Gordonias larger. Trees extend to within a few feet of the river, the brackish marshes being very narrow here. Flora of the marshes very similar to that of the Satilla River marshes near Woodbine, 16 miles north of here, but Taxodium distichum much more abundant on the St. Mary's. Crossed drawbridge about 10:50 and talked with bridge-tenders on Florida side.

25 (continued)

about 20 minutes, then came back to Georgia and collected nos. 1567-1571 in river swamp. Took two or three pictures in the vicinity.

Walked slowly back to Kingland.

Crossed Big Catfish Creek again at 2 p.m. Tide up now. Water in

Little Catfish Creek below railroad running backward, on account of tide, but absolutely nothing in the flora there to indicate proximity of the ocean. Found a fine cool spring in shady woods near Little Catfish Creek at 103d mile-post. Flora of woods surrounding spring very similar to that of similar places 200 miles inland, though this is probably not over 10 feet above sea-level. The following species were noted here:-

<i>Decumaria Barbara</i>	} climbing same tree
<i>Rhus radicans</i>	
<i>Vitis rotundifolia</i>	

25 (continued)

Magnolia grandiflora
Ilex Cassine
Asplenium Filix-foemina
Conoclinium coelestinum
Oplismenus setarius
Samolus floribundus
Potamogeton Conchites?
Aicranthemum orbiculatum
Ilex opaca
Panicum sp. (no. 1106)
Mitchella repens
Rubus nigrobaccus

Mosquitoes troublesome in these woods,
 but not seen elsewhere today.
 Met owner of spring (and adjacent
 land) a little later. He says the
 temperature of the spring is 40° F. in
 summer, and somewhat more in winter.
 Got back to Kingsland about 4:25
 and left on southbound train at
 5:47.

No Sabal Palmetto seen anywhere
 in Camden County.

- 5 (continued)

crossed St. Mary's River into Nassau Co. Fla. again at 5:53 p. m.

Very little cleared land in this part of Florida, country being nearly all flat pine-barrens (P. palustris and P. leucomensis both abundant), mostly pretty dense. Forests are partly tar-pentined but scarcely lumbered yet. Plenty of shallow Taxodium imbricarium ponds and a few Taxodium distichum swamps seen. Reached Yulee about 6:10 and had to wait there over seven hours.

Very few habitations there.

Went out with railroad night watchman between 10 and 11 p. m. to find and identify some "rattlesnake master" (Labenaria blephariglottis). The flower is supposed to bear a strong resemblance to a snake's mouth, and the crushed leaves taken internally are said to be a sure cure for snake-bite. Plant said to have several other remarkable properties. Night uncomfortably cool. Few mosquitoes around.

26. Tuesday.

Left Yulee on mixed train at 1:20 a.m., arriving at Crawford (20 miles) at 2:40. Depot closed and nobody awake. Walked up and down F. V. & W. Ry. about a mile by moonlight. Sun rose about 5:10 a.m., central time.

Changed all drivers and dined them on depot platform. Left Crawford on freight train at 10:21 a.m. Country along F. V. & W. Ry. pretty badly lumbered, both in Florida and Georgia. Crossed St. Marys River (8 miles from Crawford) back into Georgia at 11:05. More or less sand-hills on both sides of it here. Near Butler (2 miles from the river) are some fine cypress ponds, apparently untouched, and containing considerable bushy undergrowth, mostly Ilex myrtifolia. Two or three miles beyond Butler the railroad crosses a ridge (probably the same one which runs along the east side of Okefinokee Swamp) in a cut about a mile long and 6 feet deep, with a

26 (continued)

geological section exactly like that of the canal near Camp Cornelia. This ridge is doubtless what deflects the St. Mary's River so far to the southward. *Bejaria racemosa* seen several times along here. About a mile east of Moniac the country looks much like some parts of Okefinokee Swamp, the three "oaks" (*Gordonia*, *Persea*, and *Magnolia glauca*) being conspicuous, together with *Woodwardia virginica*, *Dulichium*, *Pieris nitida*, *Smilax latifolia*, etc. Got off at Moniac about 12:15 p.m.

Moniac has a temporary appearance, being strictly a sawmill town, with about 100 houses, all unpainted, mostly with 2 or 3 rooms. Walked around town some and across St. Mary's River into Baker Co. Fla. River here is only 15 or 20 feet wide, and has many logs fallen across it. *Nyssa ogeche* and *Saxodium distichum* grow in the river, and *S. imbricarium* a

26 (continued)

few feet away.

Opened trunk in freight shed (no hotel or depot here) and put yesterday's plants in press, greatly to astonishment of natives.

The following weeds have already arrived at Moniac:-

Eragrostis sp. (no. 1656)

Lyperus compressus

" *squarrosus*

Stenophyllus stenophyllus

" *Floridanus*

Euphorbia maculata.

Went down in river-swamp about 4:30 p. m. and collected nos. 1572 a-1576.

The following species were noted in addition to those mentioned above:-

Tillandsia usneoides

" sp. (no. 1573)

Cyrtia racemiflora

Ilex cassine

" *opaca*

Anastrophus paspaloides

26 (continued)

Serenoa serrulata

Lyris sp. (no. 1574)

Panicum sp. (no. 1575)

Uniola laxa

Eleocharis alnifolia

Acer rubrum.

Just outside of the river swamp are the ordinary plants of moist pine-barrens.

Jam on table at supper said to be made of fruits of *Sambucus Canadensis* and *Viburnum Ogeche*.

27. Wednesday.

Started for Okefinokee Swamp about 8 a.m. Went across river into Florida and northward along an abandoned tram road about 3 miles. Country almost uninhabited. Found Ellicott's Mound (marking east end of straight line dividing Georgia from Florida), with the assistance

27 (continued)

of a native, about 1:40 p. m. Went northwest of there about a mile without seeing anything of Okefinokee, then had to return on account of approach of night. Mound very poorly preserved and inconspicuous. Flora in the vicinity about the same as in all rather dry or moist pine-barrens in that part of the country, the following species being noted:—

*Sere noa serrulata**Ilex glabra**Eupatya flavidula**Eleocharis Baldwinii**Zypericum fasciculatum*" *opacum**Eriocaulon decangulare**Pinus serotina*" *palustris*" *bubensis**Lycopodium alopecuroides**Sarracenia minor**Sabbatia Elliottii**Centella Asiatica*

27 (continued)

Gordonia lasianthus
Taxodium imbricatum
Cratichneumon spiciformis
Drosera sp. (common)
Smilax laurifolia
Pieris nitida
Ayssa biflora
Gyrotheca tinctoria
Juncus biflorus
Trilisa odoratissima
Rhexia ciliosa
 " *glabella*
Xyris brevifolia
 " *fimbriata*
Bejania racemosa
Polygala setacea
Vaccinium nitidum ?
Marshallia graminifolia
Zyphium myrtifolium
Woodwardia Virginica
Aronia arbutifolia
Olex myrtifolia
Scleria gracilis ?
Labenaria blephariglotis

27 (continued)

Cicthra alnifolia
Orypolis filiformis
Baldwinia uniflora
Chondrophora
Physostegia
Campulocaulis aromaticus
Quercus sp. (no. 1577)
Gratiola pilosa
Asplenium cassioides
Eryngium synchaetum
Aletris farinosa
Cynoctonum sessilifolium
Dichromena latifolia
Rhynchospora corniculata
 " *axillaris*
Scleria Baldwinii
Aristida palustris
Lilium catesbaei
Polygala ramosa
Lachnocaulon anceps
Mesosphaerum rugosum
Ludwigia virgata

In the swamp of a small branch were found the following:—

27 (continued)

Nyssa Epine

Persea

Olea Virginica

Salix arundinacea

Woodwardia Virginica

Taxodium imbricatum

Eleocharis acicularis

Crossed St. Marys River near Elliott's Mound (no water in it at this time and place) at 4:34, and went down left side of it to Moniac, following an extinct and almost obliterated tram-road nearly all the way. No houses, fields, or "round timber" seen all the way back (a distance of about 4 miles). Took several pictures with both cameras on this trip. Reached Moniac at 6 p. m. Left on passenger train at 9:07, passing through about 20 miles of Baker Co. Fla. in the dark. Got off at Fargo, in Clinch Co., on right bank of Suwannee River, at 9:45.

28. Thursday

Fargo is also a sawmill town, much smaller than Moniac, but apparently built to stay, with electric lights, artesian water, painted houses, and a depot. The place is surrounded by almost unbroken pine forests. Weeds here about the same as at Moniac.

Went down to river twice in morning and took some pictures. Suwannee River here differs from all others seen, but resembles most the St. Mary's at Moniac. Banks are low but often steep, muddy or sandy, and lined with Saxodium and Nyssa Ogeche. These two species also grow in middle of river, making navigation impossible except for small boats.

Changed all diuers and put plants from Moniac in press.

Went out across river at 2:50 p.m. and collected nos. 1577b-1580 a in the swamp. Continual rain from about 3:15 to 5, making collecting disagreeable.

28 (continued)

The following species prevail in muddy flats of the river-swamp:-

Planera aquatica
Anastrophus paspaloides (no. 1585)
Sphagnum sp. (no. 1582a)
Rhexia sp. (no. 1578)
Eupatorium sp. (no. 1579)
Xyris sp. (nos 1574, 1584)
Fraxinus Caroliniana
Nyssa ogeche
Ilex myrtifolia (unusual habitat)
Scleria sp. (no. 1583)

Went beyond river a mile or so.
 Swamp on left side about $\frac{1}{2}$ mile
 wide. Dense pine-barrens beyond.
 Got back about 5:40.

29. Friday.

Cloudy all morning and threatening
 rain. Went off in woods to the eastward
 without portfolio or camera. Pine-

29 (continued)

barrens there pretty dense, timbered but not lumbered yet. Struck a tram road and followed it to Suwannee Creek and across into Echols Co. Suwannee Creek is full of Vysa Cgeche and looks much like some streams of the same size in Coffee Co. Pine-barrens on other side considerably lumbered. Collected nos. 1580a-1582 in Echols Co.

Went down on Suwannee River again in afternoon, from about 3:10 to 5:30. Collected nos. 1582a-1585 and took some $2\frac{1}{4} \times 3\frac{1}{4}$ pictures.

30. Saturday.

Went out in woods $\frac{1}{2}$ or $\frac{3}{4}$ mile north of Fargo and saw some lumbering operations with skidder, log train, etc. Changed all driers and put yesterday's plants in press. Left Fargo on freight train at

30 (continued)

1:05 p. m. Pine - barrens flat and lumbered nearly all the way to Valdosta.

Numerous tram-roads branching off from main lines, but some of them extinct. Only two or three sawmills seen, and no turpentine stills. Fine cool day.

Entered Echols Co. about 2:30. Only one curve in Echols Co., and that a very slight one, just east of Raylow.

Few cultivated fields seen until within 15 or 20 miles of Valdosta.

Crossed Allapaha River at 3 p. m.

Topography somewhat broken in the vicinity, some cuts on the railroad being 15 feet deep, mostly Lafayette.

Flora of the river banks apparently interesting but not much of it recognized. Saw Eriogonum tomentosum

near there, for first time in 17 days. Entered Lowndes Co.

about 3:25 p. m. and reached

Valdosta (47 miles from Fargo) about 3:55. Walked around the city and

30 (continued)

its outskirts some, and made notes on the flora. Topography of the vicinity heterogeneous and very puzzling. Geology probably complicated. Lafayette formation comes nearly or quite to surface in many places. Saw box of driers (missing for 17 days) in Plant System freight depot.

31. Sunday.

Started out about 10 a. m. and went down G. S. & F. Ry. a mile or two with 4x5 camera. Several cypress ponds immediately south of city, containing somewhat different flora from any seen previously. Plenty of Bastardia in them. Pine-barkens full of undergrowth. At least 6 species of Ludwigia seen.

